

وزارة
التعليم العالي والبحث العلمي
جامعة ميسان
كلية التربية الاساسية



مجلة ميسان للدراستات الأكاديمية

للعوم الانسانية والاجتماعية والتطبيقية

Misan Journal For Academic Studies
Humanits, Social and applied Sciences

ISSN (PRINT) 1994-697X

(Online)-2706-722X

المجلد 24 العدد 56 كانون الاول

vol 24 Issue 56 Dec

Misan Journal

مجلة ميسان للدراسات الأكاديمية

Misan Journal

مجلة ميسان للدراسات الأكاديمية
العلوم الإنسانية والاجتماعية والتطبيقية
كلية التربية الأساسية / جامعة ميسان

كانون الأول 2025


العدد 56

المجلد 24

DEC, 2025

SSUE 56

VOLUME 24



مركز الأبحاث في المكتبة الوطنية العراقية 1326 لسنة 2009

journal.m.academy@uomisan.edu.iq

<https://www.misan-jas.com/index.php/ojs>

<https://iasj.rdd.edu.iq/journals/journal/view/298>

الصفحة	فهرس البحوث	ت
12 – 1	Influence of the Addition of Nano Cerium Oxide/Chitosan Composite on the physical Characteristics of Polymethylmethacrylate Resin Ali Hussein Jaber Firas Abdulameer Farhan	1
27 – 13	The Impact of Peer-centred Feedback on Academic Essay Writing: A Mixed-Methods Study of Third-Year English Students at Imam Al-Kadhūm College Asmaa Hussain Jaber	2
36 – 28	Study of the evolutionary origin and virulence factors of bacterial species causing umbilical cord infections in newborns Rehab Riyadh Al-Mousawi Wafaa Abdul Wahid Al-Kaabi	3
45 - 37	Isolation and Phenotypic Characterization of Multidrug-Resistant Pseudomonas aeruginosa Isolated from Wounds and Burns of Patients in an Iraqi Clinical Setting: A Study of Their Distribution and Antibiotic Resistance Ziyad Kadhēm Dahil Alburki Samira Gjr Jremich	4
55 – 46	Genetic estimation of the toxic shock syndrome genes for burn patients in Al-Qadisiyah Province Ahmed Madboub Tahir Rana Saleh Al-Tawil	5
69 – 56	Protective effect of probiotic (Lactobacillus casei) against Escherichia coli causing diarrhea Ali J. Turki, Dhuhaa Kh. Kareem Abeer M. Alsheikly	6
84 – 70	The Impacts of Nano Barium Titanate on The Radiopacity and Surface Roughness of 3D-Printed Acrylic Denture Base Rand Naseer Kadhum Thekra Ismael Hamad	7
98 – 85	Assessment of the wettability of addition silicone Impression material following short term immersion in tea tree oil solution Samir Samier hammed Aseel Mohammed Al-Khafaji	8
110 – 99	C-peptide, liver enzymes and CRP-protein related with vitamin D deficiency in obese and diabetic (type 2) women Farah Kadhīm Alwan Ahmed Aboud khalifa	9
125 – 111	Investigation of Toxoplasma gondii in women with breast cancer by using the Histopathology technique in Southern Iraq Elaf G. G. Alzaidy Hussain A. M. Alsaady Sawsan S. Alharoon	10
139 – 126	Mapping of Gross Heterogeneity of Mishrif Formation at West Qurna 1 Oilfield, Southern Iraq Mustafa A. Abdulhasan Amna M. Handhal	11
157 – 140	A matter between two extremes: A Study in Rational Analysis Ayad Naeem Majeed	12
173 – 158	Innovation in the Introductions of the Ibn Al-Rumi's Poems (283 AH - 896 AD) Aziz Mousa Aziz	13
188 – 174	Intertextuality in the Short-Short Stories: The Case of Ahmed Jarallah Yassin Raghad Mohammed Saeed Hassan	14
210 – 189	Holograms and Virtual Sculpture: A Study in the Physical Vanishing of Digital Sculptures Works by artist Paula Dawson (as a model) Essam Nazar Mohammad Jawad	15

224 – 211	The Concepts of Predestination and Free Will in Mu'tazilite Thought (A Methodological Study from Theological to Philosophical Issues(Najlaa Mahmood Hameed	16
242 – 225	Evaluation of the Second – Grade Mathematics Textbook According to International Standards Amal Abd.A.Abass Ramla A. Kadhem	17
261 – 243	The Concept of the Hero in Ancient Iraqi Thought Atheer Ahmad Huseen Sara Saeed Abdul Redha Ekram Fares Ghanem	18
277 – 262	Synthesis and Characterization of Some 1,4-Dihydropyridine Derivatives Substituted at Position 1 and Evaluation of Their Biological Activity Sajeda Kareem Hussein Tahseen Saddam Fandi	19
291 – 278	The Syntactic Deletion in the Poetry of Al-Raai Al-Namiri Riyadh Qasim Hassan	20
303 – 292	The Language of Grammatical Criticism in Al-Radhi's Commentary on Al-Kāfiyah: A Study in Content and Style Kadhim Jabbar Alag	21
315 – 304	Visual Integration in the Structural System of Juliette Clovis's Ceramic Works: An Analytical Study of Form and Content Rula Abdul-Ilah Alwan Al-Nuaimi	22
332 – 316	An Employment of Images and Typography as a Means of Communication on Book Covers Abbas Faisal Mushtat	23
348 – 333	The Effect of Post and Brennan Strategy in Acquiring Copper Plate Skills for the Students of the Fine Arts Abbas Mahdi Jari Ronak Abboud Jaber Hussain Muhammad Ali	24
364 – 349	The Role of Contextual Learning in Raising the Level of Academic Aspiration among Students of the Department of Art Education Wiam Nadeem Jabr Al-Alaq	25
385 – 365	Environmental Degradation of the Marshes and Its Impact on Livestock Rearing (Case Study: Hammar Marsh in Dhi Qar Province) Ibtisam Ghat'a Khaji Al-Lami	26
405 – 386	Language and Gender in Riyam wa Kafa and Papa Sartre: A Lakoffian Reading Raed Hani Obaid Bany Saad Mohammed Saadi Masoud Bavanpouri	27
422 – 406	Comprehensive analysis of observed changes in pressure systems and their impact on climatic elements over Iraq (for selected climatic stations) Hassan Ali Abdul Zahra	28
443 – 423	The Effect of a Proposed Mindfulness-Based Strategy on Developing Deep Text Comprehension Skills among First-Grade Intermediate Students in Arabic Language Subject Aqeel Rasheed Abdul-Shahid Al-Asadi	29

ISSN (Print) 1994-697X
ISSN (Online) 2706-722X

DOI:

<https://doi.org/10.54633/2333-024-056-009>

Received: 15/June/2025

Accepted: 31/Aug/2025

Published online: 30/Dec/2025



MJAS: Humanities, Social and
Applied Sciences
Publishers

The university of Misan.
College of Basic Education This
article is an open access article
distributed under the terms and
conditions of the Creative
Commons Attribution

(CC BY NC ND 4.0)

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

C-peptide, liver enzymes and CRP-protein related with vitamin D deficiency in obese and diabetic (type 2) women

Farah Kadhim Alwan¹, Ahmed Aboud khalifa²

^{1,2}Department of Biology, College of Science, University of Misan, Maysan, Iraq

Email: farahka9624@gmail.com

<https://orcid.org/0009-0009-1763-4492>

Abstract :

This study aims to investigate the relationship of vitamin D deficiency and some other biochemical parameters (C-peptide, Alanine transaminase ALT, Aspartate transaminase AST, Alkaline phosphatase ALP and C-reactive protein CRP) levels in obese and diabetic women. The whole sample included 60 women aged 30 – 40 years, divided to three groups (20 women / group) as control, obesity and diabetes groups (the participants in obesity and diabetes groups were diagnosed with vitamin D deficiency). Results revealed: C-peptide increased significantly ($p \leq 0.01$) in obesity group and not significantly in diabetes group in comparison with the control, ALT increased significantly ($p \leq 0.01$) in different groups in comparison with the control, AST increased not significantly in different groups in comparison with control, ALP increased significantly ($p \leq 0.01$) in diabetes group and significantly ($p \leq 0.05$) in obesity group in comparison with the control, CRP increased significantly ($p \leq 0.01$) in

different groups in comparison with the control. The physiological impacts of these results be discussed according to the impact of vitamin D deficiency that associated with different metabolic and inflammatory disturbances, leading to diverse detrimental influence on the biochemical profiles of the studied groups.

Key words: Vitamin D deficiency, C-peptide, CRP, obesity, diabetes.

Introduction:

Vitamin D deficiency has become a global pandemic and a public health concern that involved billions of people and has contributed to multiple health complications, even in areas that receive adequate sunlight (Aparna *et al.*, 2018; Maia *et al.*, 2019).

In addition, the deficiency may be results by inadequate both sunlight exposure and nutritional vitamin intake, disorders reducing vitamin absorption, conditions that impair vitamin conversion into active metabolites, life style habits, ethnicity and genetic polymorphisms (Vierucci *et al.*, 2013).

Furthermore, this deficiency has been associated with numerous disorders, such as rickets in children and osteomalacia /osteoporosis in adults, cardiovascular diseases, arterial hypertension, dyslipidemia, cancer, multiple sclerosis, depression, dementia, psychiatric diseases, diabetes type 2 (T2D), obesity and others, but the causative role of deficiency in many of these conditions remains unclear (Milic *et al.*, 2015; Reid and Bolland 2014).

Obesity, is one of the most important risk factors for T2DM, which causes insulin resistance and inflammation due to the increase of fat tissue in the body (Rohm *et al.*, 2022). It has been reported that vitamin D receptor (VDR) changes involved in the pathogenesis of some chronic disorders such as diabetes (Isnuwardana *et al.*, 2020), the active form of vitamin D, 1,25 (OH)2D, binds to VDR and induces genes related to glucose transport, insulin secretion (Maestro *et al.*, 2003) and cellular growth in β cells (Wolden-Kirk *et al.*, 2013), in addition, vitamin D may also have a beneficial effect on the action of insulin, either directly, by stimulating the expression of insulin receptors and thereby improving insulin responsiveness to glucose transport, or indirectly, through its role in the regulation of extracellular calcium and ensuring the influx of calcium through the cell membrane (Mitri *et al.*, 2011). Vitamin D deficiency is also linked to predisposition of diabetes and may play a role in the development of diabetes (McCarthy *et al.*, 2022), moreover, a strong relationship with insulin production and insulin resistance among T2DM patients (Rafiq and Jeppesen 2021).

Vitamin D deficiency associated with inflammatory biomarkers in patients suffering from metabolic syndrome such as obesity, furthermore, serum levels of IL-6, TNF- α , and CRP trend towards higher levels in subjects with vitamin D deficiency (Khademi *et al.*, 2022; Pott-Junior *et al.*, 2020).

The relationship between vitamin D and the liver function may be attributed to the prevalence of vitamin D receptors and local conversion of this vitamin to its active form inside the non-parenchymal liver cells (Zúñiga *et al.*, 2011; Putz-Bankuti *et al.*, 2012).

Vitamin D acts as an “immune-modulator” to prevent liver cirrhosis by suppressing fibroblast proliferation and collagen production (Artaza and Norris, 2008) and is considered a factor implicated in the regulation of homeostasis, inflammation, and liver fibrogenesis (Triantos *et al.*, 2021).

Materials and methods:

Subjects:

The current study was conducted in some health centers in Maysan province / Iraq, during December 2023 to February 2024. The whole sample included 60 women aged 30 – 40 years, divided to three groups (20 women / group) as following: control, obesity and diabetes groups, the samples have been checked medically by a specialist physicians and has been diagnosed with vitamin D deficiency for obesity and diabetes groups, vitamin D(less than 20.0 ng/mL), body mass index(BMI over 30 kg/m²) and glycated hemoglobinA1c (HbA1c levels more than 6.5). Women with chronic diseases, tumors and those whom treated with hormonal drug has been excluded.

Sample collection:

Eight to ten milliliters of venous blood samples were drawn at 9 - 11 am, using a disposable needle and plastic syringes for each woman. The blood was left at room temperature for 15 minutes for coagulation, centrifuged at 3000 rpm for 5 minutes, then serum and plasma separated and transferred for storage.

Data collection and laboratory tests:

C-peptide levels were accurately measured using a Cobas e-411 / Germany with C-peptide kit, CRP levels were accurately measured using a Mindray bs_200\ China CRP kit from China. ALT, AST and ALP levels were accurately measured using a Bio systems A-15/Spain kits from Spain. The range of C-peptide, CRP, ALT, AST, ALP is 1.1 – 4.4 ng/mL, 0 – 5 mg / L, Up to 45 U/L, Up to 40 U/L and Up to 105 U/L respectively. The body mass index (BMI) was calculated as the weight in kilograms divided by the height in meters squared.

Statistical Analysis:

The statistical analysis was performed by one-way Analysis Of Variance (ANOVA), followed by Duncan's new multiple range test at a ($p \leq 0.05$), ($p \leq 0.01$) significant level (Steel *et al.*, 1997).

Results:

The results revealed :

C- peptide:

Results revealed that C- peptide increased significantly ($p \leq 0.01$) for obesity (5.75 ± 0.65 ng/ml) in comparison with diabetes (3.05 ± 0.54 ng/ml) and control (3.005 ± 0.70 ng/ml) figure (1).




Figure (1): Changes of C-peptide levels of different study's groups.

*The values represent mean $\pm$ SD.

*Different capital letters represent significant difference between groups at $p \leq 0.01$.

*Similar capital letters represent no significant difference.

CRP

Results revealed that CRP increased significantly ($p \leq 0.01$) for obesity (6.65 ± 0.94 mg/dl) and diabetes (4.56 ± 0.75 mg/dl) in comparison with control (1.50 ± 0.31 mg/dl) figure (2).

Results revealed that CRP increased significantly ($p \leq 0.01$) for obesity in comparison with diabetes figure (2).




Figure (2): Changes of CRP levels of different study's groups.

*The values represent mean $\pm$ SD.

*Different capital letters represent significant difference between groups at $p \leq 0.01$.

*Similar capital letters represent no significant difference.

ALT

Results revealed that ALT increased significantly ($p \leq 0.01$) for diabetes (20.95 ± 2.24 U/L) and obesity (16.75 ± 2.04 U/L) in comparison with control (12 ± 1.44 U/L) figure (3).

Results revealed that ALT increased significantly ($p \leq 0.01$) for diabetes in comparison with obesity figure (3).




Figure (3): Changes of ALT levels of different study's groups.

*The values represent mean $\pm$ SD.

*Different capital letters represent significant difference between groups at $p \leq 0.01$.

*Similar capital letters represent no significant difference.

AST

Results revealed that AST increased not significantly for obesity (16.10 ± 2.40 U/L) and diabetes (15.65 ± 1.49 U/L) in comparison with control (15.05 ± 1.93 U/L) figure (4).




Figure (4): Changes of AST levels of different study's groups.

*The values represent mean $\pm$ SD.

*Different capital letters represent significant difference between groups at $p \leq 0.01$.

*Similar capital letters represent no significant difference.

ALP

Results revealed that ALP increased significantly ($p \leq 0.01$) for diabetes (129.9 ± 9.00 U/L) in comparison with obesity (88 ± 3.34 U/L) and control (83.2 ± 8.20 U/L) figure (5).

Results revealed that ALP increased significantly ($p \leq 0.05$) for obesity in comparison with control figure (5).




Figure (5): Changes of ALP enzyme levels of different study groups

*The values represent mean $\pm$ SD.

*Different capital letters represent significant difference between groups at $p \leq 0.01$.

*Similar capital letters represent no significant difference.

*Different small letters represent significant difference between groups at $p \leq 0.05$.

*Similar small letters represent no significant difference.

Discussion:

The deficiency of vitamin D might be behind high C-peptide levels due to the role of vitamin D in regulation of beta cells function and insulin secretion, moreover, the metabolic and inflammatory dysregulation of the studied groups might be contributed to elevated C-peptide levels, which affect β -cell function and subsequently higher insulin secretion and insulin resistance, furthermore, high C-peptide levels might be a marker for liver dysfunction due to high current levels of liver enzyme.

Many studies agreed with the present results, many studies showed that vitamin D deficiency associated with the changes of C-peptide levels (Liu *et al.*, 2024; Xiang *et al.*, 2024; Monapati *et al.*, 2023), moreover, Al-Qahtani and his team 2024 demonstrated that vitamin D deficiency is associated with low levels of fasting C peptide for the diabetic women due the beta cells unresponsive, furthermore, in diabetic patients sufficient levels of vitamin D act synergically with high levels C peptide to controlled the diabetes status possibly due to impact of vitamin D on glucose metabolism through the function of β -cells and insulin sensitivity

High levels of C-peptide associated with high levels of CRP, this association suggests that these biomarkers may be useful in individuals identifying of developing type 2 diabetes (Lee *et al.*, 2024; Gedeberg *et al.*, 2023)

Furthermore Akter and his team 2022 showed that high C-peptide levels associated positively with the metabolic syndrome, they mention that the high BMI was the most significant factor that influencing C-peptide levels for the development of these metabolic syndrome, moreover, elevated C-peptide levels are associated with high insulin resistance in obese women, suggesting that this elevation may be considered as a risk factor for type 2 diabetes (Can *et al.*, 2023; Gilsa *et al.*, 2024)

Elevated C-peptide levels associated with high AST and ALT levels in women with NAFLD and this elevation may be useful for screening or monitoring insulin resistance in NAFLD which reflecting impaired liver function (Huang *et al.*, 2023; Fang *et al.*, 2024).

Vitamin D deficiency might be explained this increase in CRP levels due to the anti-inflammatory role of this vitamin, in addition, these groups considered as a low-grade inflammation which might be contributed of this increase, the findings of Murugiah and his colleagues 2024 are in consistence with the presents results, they found that vitamin D deficiency showed a negative association with inflammatory markers (including CRP) in diabetic women, furthermore, Different studies demonstrated that vitamin D deficiency associated negatively with CRP levels (Holmannova *et al.*, 2025; Abdelneam *et al.*, 2024), moreover, the production of pro-inflammatory cytokines decreased after vitamin binding to its receptors in monocytes, thus, vitamin D helps in reducing the concentrations of CRP and other inflammatory markers (Colin *et al.*, 2010).

Different studies pointed that diabetic and progression of associated with several inflammatory biomarkers including CRP (Okdahl *et al.*, 2022; Ellulu and Samouda, 2022; Tang *et al.*, 2022) in addition, long-term exposure to high levels of glucose significantly enhanced the increase production of pro-inflammatory cytokines, including tumor necrosis- α , interleukin (IL)-1 β , and IL-6 (Suzuki *et al.*, 2021), furthermore, high levels of CRP might be indicated for the association between pro-inflammatory cytokines and etiology of type 2 diabetes (Bahgat and Ibrahim, 2020).

Numerous studies demonstrated that vitamin D deficiency associated with high levels of ALT, AST and ALP in obese and diabetic women (Cordeiro *et al.*, 2017 ; Matar *et al.*, 2024; Fang *et al.*, 2024) due to the adverse effects of vitamin D deficiency on human liver function (He *et al.*, 2020; Skaaby *et al.*, 2014).

Furthermore, vitamin D deficiency considered as an independent risk factor for the liver enzyme's increase and the occurrence of liver diseases in nonobese , obese and diabetic women (Hashim *et al.*, 2023; Ciardullo *et al.*, 2023 ; Rajab, 2022) , in addition , vitamin D deficiency might be potentiated the adverse effects of obesity and diabetes on liver functions via its impairment of the lipid clearance by the liver , worsening the insulin resistance impact and increasing the inflammatory cytokines mediators ... that affected all together the liver functions thereby the elevation of these enzymes (Yu-Lei *et al.*, 2024; Bima *et al.*, 2021).

The association between both the high levels of liver enzymes and CRP in women with metabolic syndrome (including obesity and diabetes) represents a strong predictor of metabolic syndrome (Ghani *et al.*, 2023; Islam *et al.*, 2020; Seo *et al.*, 2019), furthermore, the overproduction of pro-inflammatory cytokines might played an important role in the liver's damage progression thereby liver enzymes elevation (Poniachik *et al.*, 2006).

In conclusion these present changes might be to vitamin D deficiency that exacerbates the imbalance of glucose metabolism and inflammatory state in obese and diabetic women, moreover, vitamin D deficiency synergistic with the metabolic group's disorders targeted the liver which is the hepatic metabolic center as evidenced by high levels of liver enzyme, thereby decline the liver abilities.

Acknowledgments:

I would like to extend my sincere thanks to Prof. Dr. Asad Yahea for own data analysis work.

Declaration of Competing Interest:

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References:

- AbdElneam, A. I., Al-Dhubaibi, M. S., Bahaj, S. S., Mohammed, G. F., Alantry, A. K., and Atef, L. M. (2024). C-reactive protein as a novel biomarker for vitamin D deficiency in alopecia areata. *Skin Research and Technology*, 30(3), e13657. <https://doi.org/10.1111/srt.13657>.

- Akter, S., Islam, M. H., Haque, M., Tasnim, N., Hasina, H., and Akther, M. T. (2022). Association of Serum C-Peptide with the Components of Metabolic Syndrome among Non Diabetic Obese. *Journal of Chittagong Medical College Teachers' Association*, 33(1), 148-153. <https://doi.org/10.3329/jcmcta.v33i1.67411>.
- Al-Qahtani, F. S., Alshaikh, A. A., and Alfaifi, S. H. (2024). The Association Between Vitamin D Deficiency and the Level of Fasting C Peptide Among Patients With Uncontrolled Type 2 Diabetes Mellitus: A Retrospective Cohort Study. *Cureus*, 16(4). DOI: 10.7759/cureus.58133.
- Aparna, P., Muthathal, S., Nongkynrih, B., and Gupta, S. K. (2018). Vitamin D deficiency in India. *Journal of family medicine and primary care*, 7(2), 324-330. DOI: 10.4103/jfmpe.jfmpe_78_18.
- Artaza, J. N., and Norris, K. C. (2008). Vitamin D reduces the expression of collagen and key profibrotic factors by inducing an antifibrotic phenotype in mesenchymal multipotent cells. *The Journal of endocrinology*, 200(2), 207. doi: [10.1677/JOE-08-0241](https://doi.org/10.1677/JOE-08-0241).
- Bahgat, M., and Ibrahim, D. (2020). Proinflammatory cytokine polarization in type 2 diabetes. *Central European Journal of Immunology*, 45(2), 170-175. <https://doi.org/10.5114/ceji.2020.97904>.
- Bima, A., Eldakhakhny, B., Nuwaylati, D., Alnami, A., Ajabnoor, M., and Elsamanoudy, A. (2021). The interplay of vitamin D deficiency and cellular senescence in the pathogenesis of obesity-related co-morbidities. *Nutrients*, 13(11), 4127. doi: 10.3390/nu13114127.
- Can, B., Ozturk, S., Gungor, K., and Sargin, M. (2023). C-peptide as an inflammatory marker in obese women. *Acta Endocrinologica (Bucharest)*, 19(1), 31. doi: 10.4183/aeb.2023.31.
- Ciardullo, S., Muraca, E., Cannistraci, R., Perra, S., Lattuada, G., and Perseghin, G. (2023). Low 25 (OH) vitamin D levels are associated with increased prevalence of nonalcoholic fatty liver disease and significant liver fibrosis. *Diabetes/Metabolism Research and Reviews*, 39(5), e3628. doi: 10.1002/dmrr.3628.
- Colin, E. M., Asmawidjaja, P. S., van Hamburg, J. P., Mus, A. M. C., van Driel, M., Hazes, J. M. W., van Leeuwen, J. P. T. M., and Lubberts, E. (2010). 1, 25-dihydroxyvitamin D3 modulates Th17 polarization and interleukin-22 expression by memory T cells from patients with early rheumatoid arthritis. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology*, 62(1), 132-142. doi: 10.1002/art.25043.
- Cordeiro, A., Pereira, S., Saboya, C. J., and Ramalho, A. (2017). Relationship between nonalcoholic fatty liver disease and vitamin D nutritional status in extreme obesity. *Canadian Journal of Gastroenterology and Hepatology*, 2017(1), 9456897. doi: 10.1155/2017/9456897.
- Ellulu, M. S., and Samouda, H. (2022). Clinical and biological risk factors associated with inflammation in patients with type 2 diabetes mellitus. *BMC Endocrine Disorders*, 22(1), 16. doi: 10.1186/s12902-021-00925-0.
- Fang, J. X., Han, Y., Meng, J., Zou, H. M., Hu, X., Han, Y. X., Huang, F., Gu, Q., and Wang, S. J. (2024). Relationship between non-alcoholic fatty liver and progressive fibrosis and serum

- 25-hydroxy vitamin D in patients with type 2 diabetes mellitus. *BMC Endocrine Disorders*, 24(1), 108. doi: 10.1186/s12902-024-01640-2.
- Gedebjerg, A., Bjerre, M., Kjaergaard, A. D., Nielsen, J. S., Rungby, J., Brandslund, I., Maeng, M., Beck-Nielsen, H., Vaag, A., Sørensen, H. T., Hansen, T. K., and Thomsen, R. W. (2023). CRP, C-peptide, and risk of first-time cardiovascular events and mortality in early type 2 diabetes: a Danish cohort study. *Diabetes Care*, 46(5), 1037-1045. doi: 10.2337/dc22-1353.
- Ghani, R., Emad, S., Khan, S. P., Naseeb, U., Ahmed, F., and Zia, S. (2023). Liver Enzyme Correlated with C-Reactive Protein as a Biomarker of Metabolic Syndrome in Elderly Patients: Metabolic Syndrome Related with Liver Enzyme and CRP. *Pakistan Journal of Health Sciences*, 280-284. DOI: <https://doi.org/10.54393/pjhs.v4i05.364>
- Gilsa, E. S., Lonappan, L., Lavanya Madhavan, S. K., and Damodaran, G. (2024). C-Peptide Levels: Correlation with anthropometric measurements of obesity and components of metabolic syndrome. *Int J Acad Med Pharm*, 6(1), 1783-1787. DOI: 10.47009/jamp.2024.6.1.352.
- Hashim, R. D., Nathir, I., Ghazy, E., Al-Timimi, I., Hameed, T. M., Hussein, M., and Ali, M. (2023). Role of plasma calcium, phosphorus, alkaline phosphatase, and parathyroid hormone in the prediction of vitamin D deficiency. *Journal of Advanced Biotechnology and Experimental Therapeutics*, 6(2), 403-411. <https://doi.org/10.5455/jabet.2023.d135>.
- He, X., Xu, C., Lu, Z. H., Fang, X. Z., Tan, J., and Song, Y. (2020). Low serum 25-hydroxyvitamin D levels are associated with liver injury markers in the US adult population. *Public Health Nutrition*, 23(16), 2915-2922. <https://doi.org/10.5455/jabet.2023.d135>.
- Holmannova, D., Borsky, P., Kremlacek, J., Krejsek, J., Hodacova, L., Cizkova, A., Fiala, Z., and Borska, L. (2025). High prevalence of low vitamin D status in the Czech Republic: a retrospective study of 119,925 participants. *European Journal of Clinical Nutrition*, 1-12. <https://doi.org/10.1038/s41430-025-01587-0>.
- Huang, J., Gao, T., Zhang, H., and Wang, X. (2023). Association of obesity profiles and metabolic health status with liver injury among US adult population in NHANES 1999–2016. *Scientific Reports*, 13(1), 15958. <https://doi.org/10.1038/s41598-023-43028-7>.
- Islam, S., Rahman, S., Haque, T., Sumon, A. H., Ahmed, A. M., and Ali, N. (2020). Prevalence of elevated liver enzymes and its association with type 2 diabetes: A cross-sectional study in Bangladeshi adults. *Endocrinology, diabetes & metabolism*, 3(2), e00116. <https://doi.org/10.1002/edm2.116>.
- Isnuwardana, R., Bijukchhe, S., Thadanipon, K., Ingsathit, A., and Thakkinian, A. (2020). Association between vitamin D and uric acid in adults: a systematic review and meta-analysis. *Hormone and Metabolic Research*, 52(10), 732-741. DOI: 10.1055/a-1240-5850.

- Khademi, Z., Hamed-Shahraki, S., and Amirkhizi, F. (2022). Vitamin D insufficiency is associated with inflammation and deregulation of adipokines in patients with metabolic syndrome. *BMC Endocrine Disorders*, 22(1), 223. doi: 10.1186/s12902-022-01141-0.
- Lee, D. H., Jin, Q., Shi, N., Wang, F., Bever, A. M., Liang, L., Hu, F. B., Song, M., Zeleznik, O. Z., Zhang, X., Joshi, A., Wu, K., Jeon, J. Y., Meyerhardt, J. A., Chan, A. T., Eliassen, A. H., Clish, C., Clinton, S. K., Giovannucci, E. L., Li, J., and Tabung, F. K. (2024). The metabolic potential of inflammatory and insulinaemic dietary patterns and risk of type 2 diabetes. *Diabetologia*, 67(1), 88-101. <https://doi.org/10.1007/s00125-023-06021-3>.
- Liu, Y. J., Duan, J. W., Lu, D. H., Zhang, F., and Liu, H. L. (2024). Association between vitamin D status and cardiometabolic risk factors in adults with type 2 diabetes in Shenzhen, China. *Frontiers in Endocrinology*, 15, 1346605. doi: 10.3389/fendo.2024.1346605.
- Maestro, B., Dávila, N., Carranza, M. C., and Calle, C. (2003). Identification of a Vitamin D response element in the human insulin receptor gene promoter. *The Journal of steroid biochemistry and molecular biology*, 84(2-3), 223-230. [https://doi.org/10.1016/S0960-0760\(03\)00032-3](https://doi.org/10.1016/S0960-0760(03)00032-3).
- Maia, F. A., Oliveira, L. M. M., Almeida, M. T. C., Alves, M. R., Saeger, V. S. D. A., Silva, V. B. D., Oliveiraa, V. S. D. D., Maria, H. M. J., Britoa, F. S. F., and Silveira, M. F. D. (2019). Autism spectrum disorder and postnatal factors: a case-control study in Brazil. *Revista Paulista de Pediatria*, 37(4), 398-405. <https://doi.org/10.1590/1984-0462/2019;37;4;00006>.
- Matar, A., Al-Tamimi, S., and Borjac, J. (2024). Evaluation of the effect of Vitamin D deficiency on liver and kidney functions of Iraqi Type II diabetic patients. *BAU Journal-Science and Technology*, 6(1), 8. DOI: <https://doi.org/10.54729/2959-331X.1139>.
- McCarthy, K., Laird, E., O'Halloran, A. M., Walsh, C., Healy, M., Fitzpatrick, A. L., Walsh, J. B., Hernandez, B., Fallon, P., Molloy, A. M., and Kenny, R. A. (2022). Association between vitamin D deficiency and the risk of prevalent type 2 diabetes and incident prediabetes: A prospective cohort study using data from The Irish Longitudinal Study on Ageing (TILDA). *EClinicalMedicine*, 53. <https://doi.org/10.1016/j.eclinm.2022.101654>.
- Milic, S., Mikolasevic, I., Krznaric-Zrnic, I., Stanic, M., Poropat, G., Stimac, D., Vlahovic-Palcevski, V., and Orlic, L. (2015). Nonalcoholic steatohepatitis: emerging targeted therapies to optimize treatment options. *Drug Design, Development and Therapy*, 4835-4845. <https://doi.org/10.2147/DDDT.S64877>.
- Mitri, J., Muraru, M. D., and Pittas, A. G. (2011). Vitamin D and type 2 diabetes: a systematic review. *European journal of clinical nutrition*, 65(9), 1005-1015. <https://doi.org/10.1038/ejcn.2011.118>.
- Monapati, S., Kaki, P., Gurajapu, M. S., Subhas, P. G., and Kudipudi, H. B. (2023). The effects of vitamin D on preventing hyperglycemia and a novel approach to its treatment. *Drugs and Drug Candidates*, 2(4), 923-936. <https://doi.org/10.3390/ddc2040046>.

- Murugiah, V., Pal, P., Sahoo, J., Nanda, N., Shamanna, S. B., and Shamanna, S. (2024). Association of Low Vitamin D Status With Adiponectin and Fibroblast Growth Factor-21 in Newly Diagnosed Type 2 Diabetes Mellitus Patients. *Cureus*, 16(10).DOI: 10.7759/cureus.71448.
- Okdahl, T., Wegeberg, A. M., Pociot, F., Brock, B., Størling, J., and Brock, C. (2022). Low-grade inflammation in type 2 diabetes: a cross-sectional study from a Danish diabetes outpatient clinic. *BMJ open*, 12(12), e062188.<https://doi.org/10.1136/bmjopen-2022-062188>.
- Poniachik, J., Csendes, A., Díaz, J. C., Rojas, J., Burdiles, P., Maluenda, F., Smok, G., Rodrigo, R., and Videla, L. A. (2006). Increased production of IL-1 α and TNF- α in lipopolysaccharide-stimulated blood from obese patients with non-alcoholic fatty liver disease. *Cytokine*, 33(5), 252-257.<https://doi.org/10.1016/j.cyto.2006.02.006>.
- Pott-Junior, H., Nascimento, C. M. C., Costa-Guarisco, L. P., Gomes, G. A. D. O., Gramani-Say, K., Orlandi, F. D. S., Gratão, A.C.M., Orlandi, A.A.D.S., Pavarini, S.C.I., Vasilceac, F.A., Zazzetta, M.S., and Cominetti, M. R. (2020). Vitamin D deficient older adults are more prone to have metabolic syndrome, but not to a greater number of metabolic syndrome parameters. *Nutrients*, 12(3), 748.<https://doi.org/10.3390/nu12030748>.
- Putz-Bankuti, C., Pilz, S., Stojakovic, T., Scharnagl, H., Pieber, T. R., Trauner, M., Obermayer-Pietsch, B., and Stauber, R. E. (2012). Association of 25-hydroxyvitamin D levels with liver dysfunction and mortality in chronic liver disease. *Liver International*, 32(5), 845-851.
<https://doi.org/10.1111/j.1478-3231.2011.02735.x>.
- Rafiq, S., and Jeppesen, P. B. (2021). Vitamin D deficiency is inversely associated with homeostatic model assessment of insulin resistance. *Nutrients*, 13(12), 4358.<https://doi.org/10.3390/nu13124358>.
- Rajab, H. A. (2022). The effect of vitamin D level on parathyroid hormone and alkaline phosphatase. *Diagnostics*, 12(11), 2828.<https://doi.org/10.3390/diagnostics12112828>.
- Reid, I. R., and Bolland, M. J. (2014). Skeletal and nonskeletal effects of vitamin D: is vitamin D a tonic for bone and other tissues?. *Osteoporosis international*, 25, 2347-2357.<https://doi.org/10.1007/s00198-014-2749-7>.
- Rohm, T. V., Meier, D. T., Olefsky, J. M., and Donath, M. Y. (2022). Inflammation in obesity, diabetes, and related disorders. *Immunity*, 55(1), 31-55.doi: 10.1016/j.immuni.2021.12.013.
- Seo, M. S., Shim, J. Y., and Lee, Y. J. (2019). Relationship between serum alkaline phosphatase level, C-reactive protein and leukocyte counts in adults aged 60 years or older. *Scandinavian Journal of Clinical and Laboratory Investigation*, 79(4), 233-237. doi: 10.1080/00365513.2019.1585567.
- Skaaby, T., Husemoen, L. L. N., Borglykke, A., Jørgensen, T., Thuesen, B. H., Pisinger, C., Schmidt, L.E., and Linneberg, A. (2014). Vitamin D status, liver enzymes, and incident liver disease and mortality: a general population study. *Endocrine*, 47, 213-220.doi: 10.1007/s12020-013-0107-8.

- Steel, R.G.D., Torrie, J.H. and Dicky, D.A. (1997). Principles and Procedures of Statistics, A Biometrical Approach. 3rd Edition, McGraw Hill, Inc. Book Co., New York, 352-358.
- Suzuki, T., Yamashita, S., Hattori, K., Matsuda, N., and Hattori, Y. (2021). Impact of a long-term high-glucose environment on pro-inflammatory responses in macrophages stimulated with lipopolysaccharide. *Naunyn-Schmiedeberg's Archives of Pharmacology*, 394(10), 2129-2139. DOI: [10.1007/s00210-021-02137-8](https://doi.org/10.1007/s00210-021-02137-8)
- Tang, M., Cao, H., Wei, X. H., Zhen, Q., Liu, F., Wang, Y. F., Fan, N.G., and Peng, Y. D. (2022). Association between high-sensitivity c-reactive protein and diabetic kidney disease in patients with type 2 diabetes mellitus. *Frontiers in endocrinology*, 13, 885516. doi: 10.3389/fendo.2022.885516.
- Triantos, C., Aggeletopoulou, I., Thomopoulos, K., and Mouzaki, A. (2021). Vitamin D—liver disease association: Biological basis and mechanisms of action. *Hepatology*, 74(2), 1065-1073. doi: 10.1002/hep.31699.
- Vierucci, F., Del Pistoia, M., Fanos, M., Gori, M., Carlone, G., Erba, P., Massimetti, G., Federico, G., and Saggese, G. (2013). Vitamin D status and predictors of hypovitaminosis D in Italian children and adolescents: a cross-sectional study. *European journal of pediatrics*, 172, 1607-1617. doi: 10.1007/s00431-013-2119-z.
- Wolden-Kirk, H., Overbergh, L., Gysemans, C., Brusgaard, K., Naamane, N., Van Lommel, L., Schuit, F., Eizirik, D.L., Christesen, H., and Mathieu, C. (2013). Unraveling the effects of 1, 25 (OH) 2D3 on global gene expression in pancreatic islets. *The Journal of steroid biochemistry and molecular biology*, 136, 68-79. doi: 10.1016/j.jsbmb.2012.10.017.
- Xiang, Q., Xu, H., Liu, Y., and Huang, W. (2024). Elevated TyG index is associated with increased risk of vitamin D deficiency among elderly patients with type 2 diabetes. *Scientific Reports*, 14(1), 16098. doi: 10.1038/s41598-024-67127-1.
- Yu-Lei, Q. U., Ying-Hao, S. O. N. G., Ru-Ru, S. U. N., and Ying-Jie, M. A. (2024). Serum Vitamin D Level in Overweight Individuals and Its Correlation With the Incidence of Non-alcoholic Fatty Liver Disease. *Physiological Research*, 73(2), 265. doi: [10.33549/physiolres.935164](https://doi.org/10.33549/physiolres.935164).
- Zúñiga, S., Firrincieli, D., Housset, C., and Chignard, N. (2011). Vitamin D and the vitamin D receptor in liver pathophysiology. *Clinics and research in hepatology and gastroenterology*, 35(4), 295-302. doi: 10.1016/j.clinre.2011.02.003.