



## Assessment of Some Heavy Metals in the Soil from Kut Al-Fadagh and Hamdan at Basrah City

Mohammed Osamah Dawood<sup>1</sup> Abdulminam H. Ali<sup>1\*</sup> Abdulkareem M. Abd<sup>2</sup>

<sup>1</sup> Department of Ecology, College of Science, University of Basrah

<sup>2</sup> Department of Biology; College of Education for Pure Sciences, University of Basrah  
E-mail:abdul\_minam.ali@uobasrah.edu.iq

### Abstract

Heavy metal accumulation in agricultural soils degrades soil quality and poses risks to crops and human health. This study aimed to measure the concentration of heavy elements in the agricultural soils of Abu Al-Khasib in the Basrah Governorate. Two sites were selected, the first site is Kut Al-Fadagh and the second site is Hamdan. Their study showed that the concentration of cobalt ranged between (21.77, ND  $\mu\text{g. g}^{-1}$ ). On the other hand, the concentration of cadmium was (1.700,0.052  $\mu\text{g.g}^{-1}$ ) while lead the values were (38.24,3.85  $\mu\text{g.g}^{-1}$ ), iron values ranged (11448,463  $\mu\text{g.g}^{-1}$ ), zinc values were (117.9,19.5), manganese values were (651.5,248.8  $\mu\text{g.g}^{-1}$ ), selenium values ranged (0.4491,0.1011  $\mu\text{g.g}^{-1}$ ) and the concentration of heavy elements in the soil varied according to the seasons. As for the average sites, Hamdan's site recorded a higher concentration of cobalt, cadmium, zinc, manganese, and selenium (0.2418,494.4,71.5,0.540,11.52  $\mu\text{g. g}^{-1}$ ) than Kut Al-Fadagh's site (0.2232,483.3,65.7,0.402,4.32  $\mu\text{g. g}^{-1}$ ). Moreover, Kut Al-Fadagh's site recorded a higher concentration of iron and lead (28.69,3467  $\mu\text{g.g}^{-1}$ ) compared to Hamdan's site, which recorded.(20.20,3067)

Keywords: Accumulation, heavy metals, pollution, soil

Received 1 /3/2023

Accepted 12 / 5/ 2023

Published 1/ 9/2024

### Introduction

Food production is necessary for life, and soil is a critical component of the natural environment that promotes plant growth. Since it supports plants, soil, along with water and air, is the basic foundation for agricultural productivity and the continuation of human life (Al-Khatib, 1998).

To conduct optimal exploitation of any natural resources and put in place appropriate management of pollution, it is necessary to identify the nature of pollutants and implement proper management and disposal strategies. Such

approaches help direct the use of priceless resources and protect them from deterioration (Youssef, 2005).

Soil contamination poses a serious threat to human health, as it can cause physiological abnormalities in the human body and its organs, which can ultimately lead to the development of deadly diseases. (Zaghloul, 2019).

Soil is exposed to a wide range of pollutants each year, either intentionally or unintentionally. Soil contamination can be classified according to its origin—either natural or anthropogenic—and according to

its nature as biological, chemical, or radioactive. Common sources of soil contamination include acid rain, radiation,

Natural weathering, such as erosion of parent rocks, atmospheric deposition, volcanic activities, etc. And anthropogenic activities such as wastewater irrigation, addition of manure, fertilizers, pesticides, are responsible for soil and crop contamination with heavy metals (HMs) (Khan *et al.*,2013).

However, one of the most significant sources of heavy metal contamination in field soil is the applied of the chemical fertilizers to promote plant growth, particularly when it used against internationally accepted standards. Additional sources include the excessive use of pesticides, combustion by-products such as charcoal and crop residues, and the application of sewage sludge from water treatment plants (Sadiq and Yousri, 2010). Heavy metals are considered very hazardous pollutants to all organisms because of their harmful impacts on living things and long-term environmental


and both domestic and industrial solid and liquid wastes (Al-Yasiri, 2016).

persistence (Cai *et al.*,2012). Therefore, the aim of the current study is to assess and evaluate the heavy elements in the field soils of two orchard regions—Kut Al-Fadagh and Hamdan—in Abu Al-Khaseeb, located in the southeastern part of Al-Basrah Governorate.

**Materials and Methods**

**Description of The Study Area**

The study area is located in Abu Al-Khaseeb, 10 km southeast of Basrah city center. This region has relatively warm winters and scorching, dry summers. Abu Al-Khaseeb was previously well known for its extensive date palm groves and numerous water channels branching from the Shatt Al-Arab River. Soil samples were collected from well-maintained and productive orchards' locations in Kut Al-Fadagh (Site 1) and Hamdan (Site 2) (Fig. 1).



**Fig.1: Study stations.**

**Soil Sample Collection** : Samples of the soil were gathered periodically between August 2022 and April 2023 from the study area using an auger. About 1.0 kilogram of soil was gathered from each location and placed in sterile plastic bags, and labeled with the necessary sample data. The soil samples were air-dried at room temperature before being sieved through a 2 mm mesh screen. Soil samples were kept dry until the conduct the further examinations.

**Soil digestion:** Soil samples were ground gently to conserve particle integrity. Followed, sieved using a 2  $\mu\text{m}$  mesh. Digestion was performed according to standard procedures described by Estefan (2013). 1 g of soil was placed into digestion tubes, and 3 mL of concentrated nitric acid ( $\text{HNO}_3$ ) was added. The tubes were heated on a hot plate at 145  $^\circ\text{C}$  for one hour. Subsequently, 4 mL of concentrated hydrochloric acid (HCl) was added while increasing the temperature. The samples were filtered using fine filter with a pore size of 0.45  $\mu\text{m}$ , and the final volume was made up to 50 mL with deionized water. Samples transferred to the Marine Chemistry Laboratory at the Marine Science Center for the determination of heavy metals using a flame Atomic Absorption Spectrophotometer (Shimadzu AA7000, Japan).

### Statistical analysis

Statistical analysis was implemented using GenStat V.7 software, utilizing a randomized complete block design (RCBD) with two factors: site and season. Differences were considered statistically significant at a probability level of  $p < 0.05$ .

### Results

Table 1 presents the outcomes of the statistical analysis of seasonal changes in heavy metal concentrations at two study sites. With regard to all different seasons and sites, the spring season at the second site (Hamdan) had the highest significant

concentration of cobalt (Co), at 21.77  $\mu\text{g}\cdot\text{g}^{-1}$ . On the other hand, the fall season at both locations had the lowest cobalt concentration (0.00  $\mu\text{g}\cdot\text{g}^{-1}$ ), which was not significantly different from the spring season at the first site (Kut Al-Fadagh).

For cadmium (Cd), the highest significant concentrations were recorded during the winter season at both sites, with values of 1.150 and 1.700  $\mu\text{g}\cdot\text{g}^{-1}$  at Kut Al-Fadagh and Hamdan, respectively. On the other hand, the lowest concentrations were recorded during the summer season at the same sites, with values of 0.052 and 0.060  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively. These low values did not differ significantly from those recorded in the autumn and spring seasons at both sites. At both study locations, these low values were not much different from those observed in the fall and spring.

Regarding lead (Pb), the highest significant value 53.79  $\mu\text{g}\cdot\text{g}^{-1}$  was recorded in the fall season at the first Site. While the lowest concentration was recorded during the same season at the same site (3.85  $\mu\text{g}\cdot\text{g}^{-1}$ ). In comparison with the first site, Hamadan recorded the highest concentrations of (Pb) in the spring season with a value of (38.24  $\mu\text{g}\cdot\text{g}^{-1}$ ) and the lowest at autumn season (4.89  $\mu\text{g}\cdot\text{g}^{-1}$ ).

For iron (Fe), the highest significant observation was recorded in the summer season at the first and second sites with a value of (11448 and 9669  $\mu\text{g}\cdot\text{g}^{-1}$ ) respectively, while the lowest value occurred in the winter at both sites (463 and 475  $\mu\text{g}\cdot\text{g}^{-1}$ ) successively.

With regard to zinc (Zn), the spring season at both sites (1 and 2) recorded the highest significant values, with 117.9  $\mu\text{g}\cdot\text{g}^{-1}$  at the second site and 109.5  $\mu\text{g}\cdot\text{g}^{-1}$  at the first site, showing no significant difference between them. Compared to the other seasons and sites, the second site's autumnal zinc concentration was noticeably lower (19.5  $\mu\text{g}\cdot\text{g}^{-1}$ ).

For manganese (Mn), the greatest value was found at the Kut Al-Fadagh and Hamdan sites ( $651.5 \mu\text{g}\cdot\text{g}^{-1}$  and  $631$ ) respectively, While The lower Mn concentration was identified in both sites in the winter season with a value ( $248.8 \mu\text{g}\cdot\text{g}^{-1}$  and  $320.2$ ).

In terms of selenium (Se), there was little variation between the two sites'

wintertime values, which were the greatest at  $0.449 \mu\text{g}\cdot\text{g}^{-1}$  at the first site and  $0.439 \mu\text{g}\cdot\text{g}^{-1}$  at the second. With no discernible difference between the two sites, the autumn season showed the lowest Se values ( $0.1100 \mu\text{g}\cdot\text{g}^{-1}$  at the first site and  $0.1011 \mu\text{g}\cdot\text{g}^{-1}$  at the second site).

**Table (1) Seasonal variations of heavy Metals ( $\mu\text{g}\cdot\text{g}^{-1}$  Dry wight) in soil**

Study Area	Seasons	Co	Cd	Pb	Fe	Zn	Mn	Se
Kut Al-Fadagh (First Site)	Summer	7.24	0.052	18.89	11448	91.0	631.0	0.1856
	Autumn	0.00	0.235	53.78	1052	34.4	460.0	0.1100
	Winter	10.05	1.150	3.85	463	28.1	248.8	0.4491
	Spring	0.00	0.172	38.23	904	109.5	593.4	0.1481
Hamdan (Second Site)	Summer	7.20	0.060	24.76	9669	94.9	651.5	0.3040
	Autumn	0.00	0.220	4.89	1226	19.5	422.0	0.1011
	Winter	17.10	1.700	12.90	475	53.5	320.2	0.4390
	Spring	21.77	0.181	38.24	898	117.9	583.8	0.1231
LSD	0.6877	0.687	3.207	370.2	8.90	10.78	0.04561	

Table 2 presents the average levels of heavy elements at the two study sites.

Hamdan site recorded the highest significant level of cobalt (Co) at  $11.52 \mu\text{g}\cdot\text{g}^{-1}$ , compared to  $4.32 \mu\text{g}\cdot\text{g}^{-1}$  at Kut Al-Fadagh site .

For cadmium (Cd) concentrations, no significant variations were recorded between the two sites, with values of  $0.540 \mu\text{g}\cdot\text{g}^{-1}$  and  $0.402 \mu\text{g}\cdot\text{g}^{-1}$ , respectively.

For lead (Pb), highest concentration at  $28.69 \mu\text{g}\cdot\text{g}^{-1}$  was observed in first site, compared to  $20.20 \mu\text{g}\cdot\text{g}^{-1}$  at the second site. In the same way, iron (Fe) recorded greater

value at the first and second sites ( $3467 \mu\text{g}\cdot\text{g}^{-1}$ ,  $3067 \mu\text{g}\cdot\text{g}^{-1}$ ) respectively .

In terms of zinc (Zn) and manganese (Mn), the second site recorded significantly higher concentrations— $71.5 \mu\text{g}\cdot\text{g}^{-1}$  for Zn and  $494.4 \mu\text{g}\cdot\text{g}^{-1}$  for Mn—compared to the first site, which recorded  $65.7 \mu\text{g}\cdot\text{g}^{-1}$  and  $483.3 \mu\text{g}\cdot\text{g}^{-1}$ , respectively. These differences between the two sites were statistically significant.

Ultimately, the selenium (Se) levels did not differ considerably between the two sites, with values of  $0.2418 \mu\text{g}\cdot\text{g}^{-1}$  at the second site and  $0.2232 \mu\text{g}\cdot\text{g}^{-1}$  at the first site.

**Table(2)The Mean accumulation Heavy metals( $\mu\text{g}\cdot\text{g}^{-1}$  Dry wight) in soil in the soil**

Study Area	Co	Cd	Pb	Fe	Zn	Mn	Se
Kut Al-Fadagh	4.32	0.402	28.69	3467	65.7	483.3	0.2232
Hamdan	11.52	0.540	20.20	3067	71.5	494.4	0.2418
<b>LSD</b>	<b>1.252</b>	<b>0.3438</b>	<b>1.603</b>	<b>185.1</b>	<b>4.45</b>	<b>5.39</b>	<b>0.02280</b>

## Discussion

### Cobalt (Co)

The elevated levels of cobalt during the winter and spring seasons may be attributed to the release of fertilizers or discharge of wastewater into irrigation channels. These results have been confirmed by Varalakshmi and Ganeshmurthy (2012),

### Cadmium (Cd)

The hike of cadmium levels in soil during the winter may be ascribed to several factors, including the use of phosphate fertilizers, the fuel combustion, and the using of contaminated irrigation water. Lower microbial activity and slower decomposition rates in winter may also reduce cadmium immobilization, allowing it to remain bioavailable in the soil. These findings are in line with those reported by Khuwaidem *et al.* (2009), who found that cadmium levels in Basrah and Baghdad exceeded world averages. Similarly, Huang *et al.* (2019) noted that the standard limit for cadmium in soil, as set by the National Environmental Quality Standards (NEQS) in China, is  $0.3 \mu\text{g}\cdot\text{g}^{-1}$ . Based on this

### Lead (Pb)

According to the National Environmental Quality Standards (NEQS), the soil quality standard for lead (Pb) contamination is  $60 \mu\text{g}\cdot\text{g}^{-1}$  (Huang *et al.*, 2019). Based on this guideline, both study sites were considered Pb-free across all seasons. These findings are in line with the results of Al-Hawi *et al.*, (2024), who

who found that fertilizer, fuel combustion, and sewage discharge are common sources of heavy metal pollution like cobalt. These results imply that agricultural fertilization activities are probably responsible for the elevated cobalt concentrations in the winter and spring.

obtained same observations within safe limits.

### Iron (Fe)

The elevation of iron concentration in the soil of both sites of present investigation during the summer season may be attributed to several factors, including workshops, factories, and vehicle discharges, as well as dust-laden sand from various sources. Additionally, some human contributions, such as the use of soil conditioners and pesticides, along with reduced river water flow during the summer months. These findings are in line with Al-Maliki and Al-Aasadi (2022), who noted similar sources of iron accumulation in soils.

The elevation of iron concentration in the soil of both sites of the present investigation during the summer season may be attributed to several factors, including workshops, factories, and vehicle discharges, as well as dust-laden sand from various sources. Additionally, some human contributions such as the use of soil conditioners and pesticides, along with reduced river water flow during the summer

months. These findings are in line with Al-Maliki and Al-Aasadi (2022), who noted similar sources of iron accumulation in soils. In the contrary, the decrease in iron (Fe) concentration during the winter season may be attributed to increased rainfall, which leads to leaching and surface runoff, causing iron and other elements to be washed out of the soil. Wuana and Okieimen (2011) indicated that the

### **Zinc (Zn)**

Current results agree with Al-Maliki and Al-Aasadi (2022), who suggested that elevated zinc levels in soil may be due to its deposition in the form of zinc sulfate ( $ZnSO_4$ ) under alkaline conditions, particularly at pH values between 7 and 9.

These results also align with Eisler (1993), who noted earlier that the main sources of zinc contamination are anthropogenic, including industrial and domestic activities, particularly from zinc-based packaging materials.

### **Manganese (Mn)**

The elevated levels of manganese (Mn) at both study sites are consistent with findings reported by Saleh (2024), who observed increased manganese concentrations in the Al-Tannumah region, east of Basrah. Similarly, Al-Bedhany et al. (2013) reported that seasonal and spatial variations significantly influence the contamination levels of various heavy metals—including iron, manganese, nickel, cobalt, copper, and lead—in the soils of Basrah. These variations are largely attributed to anthropogenic activities, industrial processes, and the geological composition of the soil. In support of this, Fosu-Mensah et al. (2017) also emphasized that the distribution and accumulation of heavy metals in soil are affected by a combination of natural factors and human activities, particularly in urban and peri-urban agricultural zones.

concentration and distribution of mineral pollutants across different regions are directly influenced by several factors, including soil properties, element transport mechanisms, and the chemical characteristics of groundwater, industrial inputs, and agricultural practices. concentration and distribution of mineral pollutants across different regions are.

### **Selenium (Se)**

According to the present findings, both the first and second sites were free from selenium (Se) contamination during all seasons, as the recorded Se concentrations did not exceed  $0.2418 \mu\text{g}\cdot\text{g}^{-1}$ . Based on the U.S. Environmental Protection Agency (US-EPA) standard, the acceptable limit for selenium in soil is  $1.6 \mu\text{g}\cdot\text{g}^{-1}$  (Saleem et al., 2012).

However, Sors et al. (2005) revealed that selenium levels in soils are influenced by various factors, including soil type, the presence of sedimentary rocks, organic matter content, and rainfall patterns. Selenium tends to accumulate more in arid and semi-arid regions, where rainfall is limited.

### **Conclusion**

This study revealed significant seasonal and spatial variations in the concentrations of heavy metals in farmland soils at Kut Al-Fadagh and Hamdan in Basrah, Iraq. While most heavy metal levels remained within internationally accepted standards, like cadmium (Cd) concentrations during the winter season exceeded the safe threshold, indicating potential environmental and public health risks. The elevated levels of certain metals were largely attributed to anthropogenic activities, including fertilizer application, waste water irrigation, and atmospheric deposition. These findings emphasize the importance of regular soil monitoring and the adoption of sustainable land management practices to prevent long-term heavy metal accumulation and ensure agricultural and ecological sustainability.

### Acknowledgement

The authors would like to express their sincere gratitude to the Marine Science Center at the University of Basrah for providing the laboratory facilities and technical support required for the heavy metal analysis. Special thanks also extend

### References

- Al- Bassam, K. S., Al- Anssari, H. R., & Khwedimm Kareem H. (2009). Study of distribution of some heavy metals in the soil of Basra City- south of Iraq. *Iraqi Journal of Science*, 50(4): 533–542.
- Al-Asadi, S. A. R., Al-Qurnawi, W. S., Al Hawash, A. B., Ghalib, H. B., & Alkhelifa, N.-H. A. (2020). Water quality and impacting factors on heavy metals levels in Shatt Al-Arab River, Basra, Iraq. *Applied Water Science*, 10(5), 103. <https://doi.org/10.1007/s13201-020-01196-1>
- Al-Baidany, A. H. (2013). Geochemical assessment of trace element in core sediments from Hor Al-Ezaim, Southern Iraq. *Marsh Bull*, 8(2): 193-202.
- Al-Khatib, A. A. (1998). Environmental chemistry of the land. Knowledge facility in Alexandria \_ Arab Republic of Egypt. P 151.
- Al Hawi, N. A., Al-Muhamed, R. A., & Al-Ali, A. K. (2024). Soil change detection and contamination assessment of some heavy metals in agricultural soils at Abu-Alkhasib, Basra Governorate, Iraq. *Iraqi Geological Journal*, 57(2F). <https://doi.org/10.46717/igj.57.2F.3ms-2024-12-13>
- Al-Maliki, H. H. N., & Al-Aasadi, W. M. T. (2022). The Assessment of Soil Contamination for Some of Basrah province Technology Regions Using ICP-GCMS. *Marsh Bulletin*, 17(2).
- Al-Yasiri, K. H. M. (2016). The effect of chemical control on soil pollution of lands located west of the city of Hilla. *Journal of University of Babylon*, 24(2): 1141–1155.
- Cai, L; Xu, Z.; Ren, M; Guo, Q; Hu, X; Hu, G; Wan, H. & Peng, P. (2012). Source identification of eight hazardous heavy metals in agricultural soils of Huizhou, Guangdong Province, China. *Ecotoxicology and Environmental Safety*, 78: 2–8. <https://doi.org/10.1016/j.ecoenv.2011.07.004>
- Eisler, R. (1993). Zinc hazards to fish, wildlife, and invertebrates: a synoptic review (Issue 26). US Department of the Interior, Fish and Wildlife Service.
- Estefan, G; Sommer, R. & Ryan, J. (2013). *Methods of Soil, Plant, and Water Analysis: A manual for the West Asia and North Africa Region* (3rd ed.). International Center for Agricultural Research in the Dry Areas (ICARDA). <https://repo.mel.cgiar.org/handle/20.500.11766/7512>
- Fosu-Mensah, B. Y., Addae, E., Yirenya-Tawiah, D., & Nyame, F. (2017). Heavy metals concentration and distribution in soils and vegetation at Korle Lagoon area in Accra. *Cogent Environmental Science*, 3(1), 1–14. <https://doi.org/10.1080/23311843.2017.1348174>
- Huang, Y; Wang, L; Wang, W; Li, T; He, Z. & Yang, X. (2019). Current status of agricultural soil pollution by heavy metals in China: A meta-analysis. *Science of The Total Environment*, 65:

- 3034–3042.  
<https://doi.org/10.1016/j.scitotenv.2018.10.185>
- Khan, K., Lu, Y; Khan, H; Ishtiaq, M; Khan, S; Waqas, M., Wei, L. & Wang, T. (2013). Heavy metals in agricultural soils and crops and their health risks in Swat District, northern Pakistan. *Food and Chemical Toxicology*, 58: 449–458. <https://doi.org/10.1016/j.fct.2013.05.014>
- Khuwaidem, K. H., Ansari, H. R., & Bassam, K. S. (2009). Study of the distribution of some heavy elements in the soil of the city of Basra in southern Iraq. *Iraqi Journal of Science*, 50(4), 542-533
- Sadiq, H., & Yousri, N. (2010). *Environmental Chemistry* (p. 303). Directorate of University Books and Publications, Publications of the University of Aleppo, Faculty of Scienc.
- Saleem, M; Essa, M. H; Chakrabarti, M. H; Low, C. T. J; Hajimolana, S. A; Hussain, M. A; Hashim, M. A, & Islam, M. S. (2012). An Economical Means for Remediating Multiple-Metal Contaminated Soil Using Electro-Kinetic Technology Under Natural Conditions. *International Journal of Electrochemical Science*, 7(9): 7809–7822. [https://doi.org/10.1016/S1452-3981\(23\)17956-9](https://doi.org/10.1016/S1452-3981(23)17956-9)
- Saleh, S. M. (2024). The concentration of heavy elements in cultivated and uncultivated soil of Basrah, Iraq. *SABRAO Journal of Breeding and Genetics*, 56(3), 1298–1307. <https://doi.org/10.54910/sabrao2024.56.3.37>
- Sors, T. G; Ellis, D. R. & Salt, D. E. (2005). Selenium uptake, translocation, assimilation and metabolic fate in plants. *Photosynthesis Research*, 86(3): 373–389. <https://doi.org/10.1007/s11120-005-5222-9>
- Varalakshmi, L. R., & Ganeshamurthy, A. N. (2012). Heavy Metal Contamination of Water Bodies, Soils and Vegetables in Peri-Urban Areas: A Case Study in Bangaluru. *Journal of Horticultural Sciences*, 7(1): 62–67. <https://doi.org/10.24154/jhs.v7i1.392>
- Wuana, R. A. & Okieimen, F. E. (2011). Heavy Metals in Contaminated Soils: A Review of Sources, Chemistry, Risks and Best Available Strategies for Remediation. *ISRN Ecology*, 2011: 1–20. <https://doi.org/10.5402/2011/402647>
- Youssef, A. F. (2005). *Soil and water analysis devices and methods* (1st ed., p. 551). King Saud University.
- Zaghloul, A. A. F. (2019). *Environmental pollution ... Problems and solutions*. Al-Huda Center for Computer and Photography Press

## تقييم بعض المعادن الثقيلة في التربة من كوت الفداغ وحمدان في مدينة البصرة

\* محمد أسامة داود \* عبد المنعم حسين علي \*\* عبد الكريم محمد عبد  
 \*قسم البيئة ، كلية العلوم ، جامعة البصرة ، العراق  
 \*\*قسم علوم الحياة ، كلية التربية للعلوم الصرفة ، جامعة البصرة ، العراق

## الملخص

تعد المعادن الثقيلة خطيرة بسبب ثباتها وسميتها. قد يؤدي التلوث بالمعادن الثقيلة في التربة الزراعية إلى اخلال في وظائف التربة، والتدخل في نمو المحاصيل، وحتى الإضرار بصحة الإنسان من خلال تلوث السلسلة الغذائية. هدفت الدراسة إلى قياس تركيز العناصر الثقيلة في الترب الزراعية لمنطقة أبي الخصيب في محافظة البصرة، وتم اختيار موقعين الموقع الأول هو كوت الفداغ والموقع الثاني حمدان. أظهرت الدراسة أن تركيز الكوبالت تراوح بين (ND. 21.77) مايكروغرام.غرام<sup>-1</sup> في المقابل بلغ تركيز الكاديوم (1.700.0.052) مايكروغرام.غرام<sup>-1</sup> فيما بلغت القيم الرصاص (38.24.4.89) مايكروغرام.غرام<sup>-1</sup> وتراوحت قيم الحديد (11448.463) مايكروغرام.غرام<sup>-1</sup> وكانت قيم الزنك (117.9.19.5) مايكروغرام.غرام<sup>-1</sup> وكانت قيم المنغنيز (651.5.248.8) مايكروغرام.غرام<sup>-1</sup>. وتراوحت قيم السيليونيوم (0.4491،0.1011) مايكروغرام.غرام<sup>-1</sup> وتباينت تراكيز العناصر الثقيلة في التربة باختلاف الفصول. أما متوسط المواقع فقد سجل موقع حمدان أعلى تركيز لعناصر الكوبالت والكاديوم والزنك والمنجنيز والسيليونيوم (0.2418,494.4,71.5,0.540,11.52) مايكروغرام.غرام<sup>-1</sup> مقارنة بموقع كوت الفداغ الذي سجل (0.402,4.32, 0.2232,483.3,65.7). كما سجل موقع كوت الفداغ تركيزاً أعلى للحديد والرصاص (28.69,3467) مايكروغرام.غرام<sup>-1</sup> مقارنة بموقع حمدان الذي سجل (20.20,3067) مايكروغرام.غرام<sup>-1</sup>