Kirkuk University Journal for Agricultural Sciences
ISSN:2958-6585
https://kujas.uokirkuk.edu.iq

RESEARCH ARTICLE
https://doi.org. 10.58928/ku26.17102

Accumulation of Lead and Cadmium inM (Zea mays L.) Plants Growing in Soil Adjacent to the Northern Gas Company in Kirkuk

Fatima Jamal Kamel[1] Sameerah Faydhallah Mohammed [1] Akram Othman Esmail[2]
[1]*Department of Soil and Water Resource, College of Agriculture,University of Kirkuk, Kirkuk, IRAQ.*
[2]*Department of Soil and Water Resource, College of Agriculture,University of Salahaddin. Erbil, IRAQ.*
*Corresponding Author: aksm24053@uokirkuk.edu.iq.

Received:09/07/2025 Revised:05/ 09/2025 Accepted: 02/01/2026 Published: 04/03/2026

ABSTRACT

The area surrounding the North Gas Company, was chosen to study the pollution resulting from the North Gas Company and its impact on plants in the surrounding area with (lead and cadmium), as well as to estimate the bioaccumulation of plants. Three sites were identified to estimate the bioaccumulation at a distance of (50, 1255 and 3150) meters. Soil samples were collected at a depth of (0-20) centimeters, September, 2024.regarding the plant , two samples were taken at the first station, one sample for each farm, while at the second and third stations Five samples per each station were taken, one sample per farm, in the form of a semicircle surrounding the company. The results showed that all sites had BCF values exceeding 1, with the exception of the third site for lead. This indicates that the plants have lead and cadmium were more concentrated in it than in the soil, and plants have a high potential to accumulate these elements (hyperaccumulator potential).

***Keywords*:** Heavy Metals, Plants, bioconcentration factor, Kirkuk City.

Copyright © 2026. This is an open-access article distributed under the Creative Commons Attribution License.

INTRODUCTION

Plant pollution is the accumulation of pollutants within plant tissues, such as toxic elements and pesticides. The response of plants to various toxic elements present in the soil varies depending on the plant species, age, and growth stage of the plants. It is also affected by the physicochemical properties of the soil, root secretions, and the concentration and type of toxic elements in the soil, in addition to the difference in solubility, absorption, transport, and chemical effectiveness of toxic elements. All of this leads to differences in toxicity to plants[1] ‹The increase in the concentrations of these elements in plants comes from the growth of the plant in soil contaminated with these elements [2]. The phenomenon of accumulation is a natural phenomenon in plants that accumulate toxic elements without showing symptoms of toxicity. There are approximately 400 plant species belonging to 45 families called accumulating plants, which are characterized by the ratio of the element's exacerbation in the vegetative system to the root system being greater than one, while for non-accumulating plants it is less than one [3]. There is a significant relationship between the concentration of heavy elements in the soil and the plants growing in it, and the difference in the concentration of any element in any type of plant is due to the difference in the active groups that are associated with the toxic elements for plants, which leads to differences in absorption efficiency and accumulation between plant species and varieties [4] that polluted soils have a negative impact on plants, but there are many plant species that have the ability to tolerate high concentrations of these elements without affecting their growth. Thus, these plants can be exploited in the process of reclamation and remediation of soils contaminated with toxic elements by absorbing and accumulating them within various plant tissues such as roots, stems, and leaves. This is known as bioremediation or phytoremediation. This technology is considered one of the promising solutions in treating soil pollution problems. It is called green technology through which the concentration of pollutants in the soil is reduced [5]. The bioaccumulation factor (BCF) is important in knowing the plants used in phytoremediation of soils contaminated with safe elements [6][7] The aim of this research is to study the effects of lead and cadmium on maize plants, and then estimate the bioconcentration factor (BCF) for the studied plants in the study area.

Materials And Methods

1- The study area and sample collection: The area surrounding the North Gas Company, which is about 24 kilometers from the city center of Kirkuk, was chosen. Three station were identified to estimate the concentration of plant biomass at a distance of (50, 1255 and 31255) meters. Soil samples were collected from the surface of the soil at a depth of (0-20) centimeters in,September,2024 and then transferred to the laboratory for the purpose of conducting physical and chemical analyses. while in regard to the plant, samples were collected from the same sites from which soil samples were collected ,

two samples were taken at the first location one sample for each farm. On the other hand, at the second and third location have Five samples per station were taken, and one sample per farm, in the form of a semicircle surrounding the company, as shown in Map (1).

Map (1) showing the location of the North Gas Company and sampling points.

2 . Soil Physical and Chemical Analysis

The study soil physical and chemical analyses were conducted in the soil laboratory at the College of Agriculture, University of Kirkuk.

2.1. Soil texture: Particle size distribution (PSD) estimated using the hydrometer method described by [8], as well as the texture triangle.

2.2. Soil moisture content was estimated using the gravimetric method as described by Jensen [9].

2.3. Bulk density was estimated using the paraffin wax method described by Black [10].

2.4. pH and Electrical conductivity (EC) estimated using a (Hanna model) pH-meter and EC-meter from a soil extract (1:2.5) according to the method described in Page [11]and EC expressed in units (ds m-1).

2.5.Organic matter (OM) ,measured using the wet oxidation method according to the Smith Weldon modification of the Walkey-Black method described in Black[10] Calcium carbonate (CaCO3) was measured according to Piper (1950) as described in [12].

3. Total heavy metal estimation: The digestion process was carried out according to the method of [13] to estimate the total heavy metals.

4. Available heavy metals: extracted using a chelating compound of 0.005 mM (DTPA), 0.01 mM (calcium chloride), and 0.1 mM (triethanolamine (TEA) in a ratio of 2:1 (soil: extraction solution) according to the method described in [14].

5. Total and available heavy metal in the extracts were estimated using the Jena atomic absorption analyticaly (AAA).

6.Digestion of plant samples and determination of heavy elements in them The digestion process was carried out by taking 0.2 of the dry sample, then 4 ml of sulfuric acid was added and left overnight. After that, 2 ml of perchloric acid was added and exposed to heat as stated by [11].

7. Estimation of plant chlorophyll

Chlorophyll was estimated using 10 ml of ethyl alcohol and 0.50 g of the plant sample according to [15].

1.0 1.mg chlorophyll a/ml solution = (13.70)(A665nm) - (5.76)(A649nm)……..1

2.0 2.mg chlorophyll b/ml solution = (25.80)(A649nm) - (7.60)(A649nm)……2

8. Estimation of Bioconcentration Factor (BCF)

The accumulation of the element in cultivated plant species was estimated using the Bioconcentration Factor method according to [16] as shown in the following equation:

$$BCF=\frac{(Concentration\ of\ the\ element\ in\ the\ plant\ )}{(Concentration\ of\ the\ element\ in\ the\ soil\ )}\ldots\ldots\ldots.3$$

9. Statistical Analysis

9.1. Analysis of Variance (ANOVA): - Significant differences in results between samples were determined based on station and seasonal differences and their overlap using the ANOVA method. Averages were compared using the LSD test at a significance level of 0.01 using SPSS Version 27.[17]

9.2. Pearson Correlation Coefficient - The correlation coefficient between the studied factors was calculated on a computer using the SPSS Special Program for Statistical Analysis (SPSS) software. This is a Pearson System Correlation Coefficient.[17].

Results And Discussion

1. Soil Physical and Chemical Analysis

1.1. Soil Texture: The soil texture of the study area was classified as medium to fine. It is well known that several factors are responsible for the soil formation and development, according to the equation [18].

1.2.Bulk density (Bd): Table (1) Shows non-significant influences of location on bulk density the highest value of 1.64 Mg m^{-3} was recorded at the second location and the lowest value of was 1.60 Mg m^{-3} at the third location , respectively. This is consistent with the findings of [19], where the Bd value before planting was 1.45 Mg m^{-3} in the soil of the College of Agriculture Research location - University of Kirkuk.

1.3.Soil Moisture Content% :Table (1) shows that the location did not have a significant effect at the level of significant 0.01 on the soil moisture content, as the highest percentage of 3.62% was recorded in the third location and the lowest percentage of 3.38% in the first location.

1.4.The ionic potential of hydrogen (pH) as shown in Table (1) that the location have a significant effect on the pH values, as the lowest value of 6.95 was recorded in the second location and the highest value of 7.44 in the third location, respectively. as the pH value before planting was 7.1 in the soil of the College of Agriculture Research Station - University of Kirkuk. It also partially agrees with what was reached by the researchers [20], The pH value was 7.72 before planting in the soil of the (Koldara) area, affiliated with the Alton Kopri district (Dibs district), Kirkuk Governorate.

1.5.Electrical conductivity (EC): It was found that had a significant effect, as the highest value of 1.58 dSm-1 was recorded at the third location and the lowest value of 1.18 dSm-1 at the first location , Table (1) "The results were in agreement with the researcher's conclusion, indicating that the electrical conductivity was 0.38 dSm^{-1} in the soil of the Grdarasha Experimental Field Research Center, College of Agricultural Engineering Science[21].[22] where the EC Value before planting was 2.3 dSm-1 in the soil of the College of Agriculture Researchers location -University of Kirkuk. [23]The EC value was 2.17 before planting in the soil of the (Koldara) area, affiliated with the Alton Kopri district (Dibs district), Kirkuk Governorate.

1.6.Organic matter (OM) Table (1) shows that the location did not have a significant effect at the 0.01 level on the organic matter of the soil, as the highest rate was 1.10% at the first site and the lowest rate was 0.96% at the second and third location ,[24] The organic matter value was 0.97 before planting in the soil of the (Koldara) area, affiliated with the Alton Kopri district (Dibs district), Kirkuk Governorate.consistent .

1.7.Calcium carbonate (CaCO3) Table (1) shows that the location have a significant impact, with the highest percentage of 31.50% recorded at the second location and the lowest percentage of 25.86% at the first location.

2. Total and available lead concentration in the soil

2.1. Total lead concentration (Pb T): Table (1) shows significant differences between Pb concentration at the first and second location, where the highest concentration (98.15 ppm) appeared at the third stations and the lowest concentration (84.34 ppm) was recorded at the first location.

2.2. Available lead concentration (Pb A): Table (1) explains that there are significant differences between available concentration of Pb between the study location at the level of significant 0.01, where the highest concentration of 16.09 ppm was recorded at the third location and the lowest concentration of 12.97 ppm was observed at the first location.

3. Total and available cadmium concentration in the soil

Total cadmium concentration (Cd T): Table (1) shows that there are significant differences between Cd concentration between the study location , the highest concentrations was noted from the first and second location with the values of (22.36 and 21.30) ppm respectively, while the lowest concentration of 17.84 ppm was recorded at the first location On the other hand, the Available cadmium concentration (Cd A) was not detected.

Table (1) Impact of location on the chemical and physical properties of the soil and concentration of total and available lead and cadmium

parameters	location 1	location 2	location 3	L.S.D0.01
EC	1.18	1.36	1.58	0. 39
pH	7.29	6.95	7.44	0.33
$CaCO_3$	25.86	31.5	26.95	4.60
O.M%	1.10	0.96	0.96	0.71N.S
Bd $Mg.m^{-1}$	1.62	1.64	1.60	0.74 N.S
M%	3.38	3.60	3.62	0.21N.S
PbT ppm	84.34	86.12	98.15	13.81
Pb A ppm	12.97	14.66	16.09	1.42
Cd T ppm	17.84	22.36	21.30	3.80
Cd A ppm	N.D.= not detected			

4. Concentration of heavy elements in the plant

4.1. Plants lead concentration: Table (2) indicates to significant differences at level of significant of 0.01 between the stations. The highest average of 487.5 ppm was recorded at the second location and the lowest average of 22.25 ppm at the third location . It is clear that all concentrations and all location exceeded the limits permitted by the World Food and Health Organization, which is 5 ppm [25]. The reason for this is that the harmful effects of this company lead to increase the pollution of nearby agricultural soils with this element, as well as increasing its absorption by plants in these soils. This is consistent with what was indicated by [26], who showed the increased absorption of heavy elements by plants that pollute the soil,

leading to their accumulation in the soil due to the heavy pollution caused by industrial facilities and factories near these lands. Also, the great ability of some plants to absorb and accumulate large quantities of lead in their biomass, reaching 2300 ppm dry matter [27].

4.2. Concentration of cadmium element in the plant (Cd P) : Table (2) shows that there are significant differences at level of significance 0.01 between Cd concentration in the study location, where the highest average of 99.702 ppm was recorded at the third stations, and the lowest average of 62.400 ppm was observed at the first station. We note an increase in cadmium concentrations over the permissible limits of the World Food and Health Organization [25], which is 0.2 ppm, and the reason is due to its high absorption by the plant and its accumulation in the plant tissues, and this results in agreement with those recorded by [28], when studying the role of some industrial facilities in Baghdad city in polluting the soil, water and plants with some heavy elements. Cadmium concentrations in the plant vary according to the type of plant and plant parts. It is higher in the roots than in the rest of the other parts of the plant, and its concentration in the herbs ranges between 50-320 ($\mu g\ g^{-1}$) of dry substance [29].

5. Chlorophyll content (mg g^{-1}): Table (2) shows that there are no significant differences at the 0.01 level between the three location , as the highest rate of 5.18 mg g^{-1} was recorded at the second location and the lowest rate of 4.90 mg g^{-1} was recorded at the first location.

Table 2: Lead, cadmium and chlorophyll content in maize plants

Sequence	Location	Samples	Chlorophyll mg g^{-1}	Cd plant ppm	Pb plant ppm
1	First location	S_1R_1	4.60	33.70	407.50
2		S_1R_2	5.21	91.10	470.00
First Location average			4.90	62.40	438.75
3	second location	S2R1	5.22	87.45	487.50
4		S2R2	5.22	94.75	102.00
5		S2R3	5.22	99.52	50.47
6		S2R4	5.12	104.60	52.82
7		S2R5	5.16	103.22	55.25
second Location average			5.18	97.91	149.61
8	Third location	S3R1	5.22	103.22	45.75
9		S3R2	5.22	102.20	22.25
10		S3R3	4.88	86.77	59.00
11		S3R4	5.19	103.97	47.50
12		S3R5	5.22	93.00	53.75
Third Location average			5.14	99.70	45.30
L.S.D 0.01			0.4 N.S	13.20	107

6.Bioconcentration Factor(BCF):

It is possible to estimate the amount of the element absorbed by the plant by estimating the bioconcentration factor as follow: The ratio between the concentration of an element in the plant and its concentration in the soil.. The results showed that the lowest bioconcentration factor (BCF) for lead absorption appeared in the maize plant, 0.62 at the third location , and the highest value was 6.41 at the first location , while the lowest value for the bioconcentration factor for cadmium was recorded in the maize plant, 5.19 at the first location , and the highest value , 6.05 at the third location , which indicates the variation in the plant's ability to take the toxic element from the soil. It was found that the bioconcentration capacity for cadmium is greater than that for lead, as shown in Table (5). Lead accumulates easily on the soil surface, and the source of contamination with this toxic element is gasoline, batteries, old grease, pesticides, and others [30]. It is shown from Table (5) that all location had BCF exceeding 1, except for the third site for lead. This indicates that the plants had concentrated lead and cadmium more than the soil, and that the plants have a hyperaccumulator potential for the element, This is consistent with study of [31], which studied the bioconcentration factor by estimating some heavy elements in soils and common plants in different habitats in Kirkuk City, It was found that most plants have the ability to accumulate high amounts of heavy metals in their leaves.

Table (5) Bioconcentration Factor (BCF) in maize plants in the study area.

location	Lead	Cadmium
First location	6.41	5.19
second location	2.40	6.32
Third location	0.62	6.05

7. Pearson Correlation Coefficient:

Table (6) shows that the concentration of lead in plants has a significant negative relationship with the concentration of cadmium in plants, as the correlation coefficient value (r)reached 0.58 *. It also has a significant negative relationship, of r= -0.33 with the total concentration of cadmium in the soil, while the concentration of cadmium in plants has a highly significant relationship with chlorophyll, as the correlation coefficient (r) reached to 0.73**. It also has a significant positive (direct) relationship with the concentration of available lead, as the correlation coefficient0.60*.

Table (6) Pearson correlation relationships between soil lead and cadmium concentrations , plants lead and cadmium concentrations and chlorophyll content.

	Pb-p	Cd-p	Chll	Pb-T	Cd-T	Pb-A
Pb-p	1					
Cd-p	-0.58*	1				
Chll	-0.28	0.73**	1			
Pb-T	0.19	-.38	-0.47	1		
Cd-T	-0.33	0.47	0.16	0.17	1	
Pb-A	-0.763**	0.60*	0.29	0.21	0.55	1

*. Correlation is significant at the 0.05 level (2-tailed).
**. Correlation is significant at the 0.01 level (2-tailed).

Conclusion

The study showed that the concentration of lead in plants for all sites exceeded the permissible limits of the World Food and Health Organization which is 5 ppm [25]. The concentration of cadmium also exceeded the permissible limits of the World Food and Health Organization [25], which is 0.2 ppm . All location had BCF value exceeding 1, except for the third location for lead. This indicates that the plants had concentrated lead and cadmium more than the soil, and that the plants have a hyperaccumulator potential for these elements.

References

[1]. DalCorso, G.; A. Manara and A. Furini.(2013). An overview of heavy metal challenge in plants: from roots to shoots. Metallomics 5: 1117–1132..

[2]. Azita, B.H. and Seid, A.M.(2008). Investigation of heavy metals uptake by vegetable crops from metal-contaminated soil. World Academy of Science, Engineering and Technology 43(1):56-58.

[3]. Al-Wahaibi, Muhammad bin Hamad (2007). The phenomenon of the presence of the basic element in plants. Saudi Journal of Life Sciences, Journal (14), Issue 2.

[4]. Hajar, E. W. I.; Sulaiman, A. Z. B., and Sakinah, A. M. (2014). Assessment of Heavy Metals Tolerance in Leaves, Stems and Flowers of Stevia rebaudiana Plant. Procedia Environmental Sciences 20: 386-393.

[5]. Lahori, A.H., Ahmad, S.R., Afzal, A., Mierzwa- Hersztek, M., Bano, S Muhammad, M.T., Saleem, I., Soomro, W.A. (2023). Alone and combined applica- tion of press mud compost and fuller earth for abating Pb and Cd and enhance sorghum growth in polluted soils. Trends in Ecological and Indoor Environment Engineering, 1(1), 7–15.

[6]. Wei, S.; Q. Zhou, and P. Koval. (2006). Flowering stage characteristics of cadmium hyperaccumulator Solanum nigrum L. and their significance to phytoremediation. Sci. Total Environ. 369(1-3): 441-446 doi:10.1016/j.scitotenv.2006.06. 014. PMID:16859734.

[7]. Zacchini, M.; F. Pietrini; G. Scarascia Mugnozza; V. lori; L. Pietrosanti, and Massacci, A. (2009). Metal tolerance, accumulation and translocation in poplar and willow clones treated with cadmium in hydroponics. Water Air Soil Pollut. 197(1-4): 23-34. doi:10.1007/s11270-008-9788-7.

[8]. Gee, G. W. and Bauder, J. W. (1986). Partial size analysis. In Methods of Soil Analysis Part(1): Physical and Mineralogical Methods (2nd ed.). Edited by A. Klute, PP: 383-409.

[9]. Jensen, J.R. (1983). Chloride dispersion in packed columns during saturated steady flow. J. Soil Sci. 34:249-262.

[10]. Black, C.A. (1965). Method of soil analysis. Am Soc. of Agro. No. 9, Part 1 and 2.

[11]. Page, A.L., R.H. Miller and Kenny, D.R. (1982). Method of soil andlysis part (2), 2nd ed. Agronomy series 9. Amer. Soc of Agron, Madison, Wisconsin. Potassium in Soil: Amini review. Cheni. Int. 2 (1), 58-69.

[12]. Hesse, P.R. (1972). A text book of Soil chemical analysis. Chemical Publishing Co, Inc. New York, pp. 240-250.

[13]. Jones, D.L.; T. Eldhuset; H.A. de Wit, and B. Swensen. (2001). Aluminum effect on organic acid mineralization in a Norway spruce forest soil. Soil Biology and Biochemistry 33(9):1259-1267.

[14]. Lindsay, W. L. and W. A. Norvell. (1978). Development of a DTPA soil test for Zinc, Iron, Manganese, and Copper 1. Soil Science Society of America Journal. 42(3):421-428.

[15]. Wintermans, J.F. and Demote, A.,)1965(. Spectrophotometry characteristics of chlorophyll (a) and (b) and their phytophytins in ethanol. Biochimica et Biophysica Acta (BBA) - Biophysics including Photosynthesis, 109, pp.448-453.

[16]. Baker AJM, Reeves RD, Hajar ASM (1994). Heavy metal accumulation and tolerance in British populations of the metallophyte Thlaspi caerulescens. New Phytol. 127: 61-68.
[17]. Esmail, A. O., Mustafa, A. O. and Omar, Q. A. 2024. Experimental design and analysis. Faculty of Agriculture - Salahuddin University - Erbil - Press of the Ministry of Agriculture and Irrigation Kurdistan Region. Iraq. Number of pages: 275 .
[18]. Jenny, H. (1941). Factors of soil formation: a system of quantitative pedology. Courier Corporation.
[19]. Alobaidy, A. S., Tahir, H. T., & Alshamary, W. F. (2024). The role of the use of some soil enhancers and the type of subsurface drip irrigation tube in some of the physical properties of the soil and the efficient use of water for the Growth and yield of eggplant crop under protected farming conditions. Journal of Kirkuk University for Agricultural Sciences, 15(4).
[20]. Sharif, Y., & Al Jubor, J. (2020). Effect of Plant Density and Organic Fertilizers on Growth and Yield of Pop Corn (Zea mays L. everta). Kirkuk University Journal For Agricultural Sciences, 11(1), 126-143. https://doi:10.58928/ku20.11111.
[21]. Abdullah, S.A., Ali, V.D., Salih,R.H.,&Muhammed,B.I.(2024).Response of Growth and Yield Traits of Safflower Varieties(Carthamus tinctorius L.)to Zinc Foliar Application .Journal of Kirkuk University for Agricultural Sciences,15(40).
[22]. Al-Shamary, W.F.A., Sharif, Y.O.N.A., Noori, N.E., and Kahlel, A.S. (2025). Effect of Moisture Depletion Rate and Irrigation Water Depth on the Productivity and Water Use Efficiency of Soybean Crop (Glycine max L.) Merr. under Drip Irrigation and Fixed Sprinkler Irrigation Systems. Agricultural Science Digest. 45(2): 222-227 https://arccjournals.com/journal/agricultural-science-digest/DF-.
[23]. Sharif, Y. O. N. A., Al-Mafrajy, A. S. H., & Albarzenje, Z. M. M. (2024). Evaluation of the Effectiveness of 2.4-D 72% Herbicide in Combating Weedy Leaves Associated with Different Varieties of Forage Sorghum (Sorghum bicolor L.) and Its Effect on Growth Characteristics. IOP Conference Series: Earth and Environmental Science, 1371(5). https://doi.org/10.1088/1755-1315/1371/5/052020.
[24]. Sharif, Y. O. N. A., Madab, D. S., & Hindi, H. A. (2024). Estimation of Path Analysis and Genetic Parameters for Sorghum (Sorghum bicolor (L.) Moench) Varieties in Different Environments. IOP Conference Series: Earth and Environmental Science, 1371(5). https://doi.org/10.1088/1755-1315/1371/5/052035.
[25]. WHO/FAO. (2007). Joint WHO/FAO. Food standard programme codex Alimentarius commission 13th session.
[26]. Azita, B.H. and Seid, A.M.)2008(. Investigation of heavy metals uptake by vegetable crops from metal-contaminated soil. World Academy of Science, Engineering and Technology 43(1):56-58.
[27]. Yang, P.; R. Mao; H. Shao and Y. Gao. (2009). The spatial variability of heavy metal distribution in the farmland of Taihang Piedmont plain, China. C. R. Biologies. 332:558-566.
[28]. AL-Ghalibi ,Dhay Mahdi Saleh.(2016).Role of some industrial installations in Baghdad city on soil,plant and water pollution by some of heavy metals . Master's Thesis. College of Agriculture, University of Baghdad.
[29]. Kabata, Pendias A. and H. Pendias.(2001). Trace elements in soils and plants. 3rd ed., CRC Press. Boca Raton, FL.
[30]. Boamponsem, G. A.; M. Kumi and I. Debrah (2012). Heavy Metals Accumulation in Cabbage, Lettuce and Carrot Irrigated with Wastewater from Nagodi Mining Site in Ghana. Intern. J. Sci. & Technol. Res. 1 (11): 124-129. Bohan, L.; Z.
[31]. Abdljaba, R,A., &Muhammad, S.F. (2017). Estimation of Metal Accumulation by Boiconcentration Factor Through The Determination of some Heavy Metals in soil and Common Plants in Different Habitats in Kirkuk City Tikrit Journal for Agricultural Sciences,17(3).

تراكم عنصري الرصاص والكادميوم في نباتات الذرة الصفراء (Zea mays L.) النامية في الترب المجاورة لشركة غاز الشمال /كركوك

فاطمة جمال كامل[1] **سميرة فيض الله محمد[1]** **اكرم عثمان اسماعيل[1]**

[1]قسم علوم التربة والموارد المائية ، كلية الزراعة , جامعة كركوك , كركوك ,العراق.

الخلاصة

تم اختيار المنطقة المحيطة بشركة غاز الشمال لدراسة تلوث الناتج من شركة غاز الشمال وتاثيرها على النباتات في المنطقة المحيطة بها ببعض العناصر الثقيلة (الرصاص والكادميوم) فضلا عن تقدير التراكم الحيوي للنبات ، تم تحديد ثلاث مواقع لتقدير التراكم الحيوي على بعد(50، 1255،3150متر)، موقع الاول نموذجين والمحطة الثانية والثالثة خمس نماذج على شكل نصف دائرة المحيطة بالشركة ، تم اخذ عينات النبات والتربة الخاصة بة خلال الموسم الاول في حين جمعت عينات التربة فقط لموسمين من منطقة المحيطة وتم اجراء تحليل الكلورفيل وعنصري الرصاص والكادميوم للنبات و اجراء التحاليل الكيميائية والفيزيائية للتربة في موسمين) 2024) و(2025) اذ جمعت عينات التربة على عمق (0-20)سم ، ، وبينت النتائج وجود فروق معنوية لكل من تركيز عنصري الرصاص والكادميوم والكلوروفيل في النبات للمحطات الثلاثة المحددة في هذه الدراسة ،ان جميع المواقع قد تجاوزت فيها قيم ***BCF*** عن 1 باستثناء الموقع الثالث للرصاص وهذا يدل على ان النباتات قد تركزت فيها عنصري الرصاص والكادميوم اكثر من التربة وان النباتات ذات امكانية تراكم فائقة للعنصري (***hyperaccamulator potential***).

الكلمات المفتاحية: العناصر الثقيلة ، نباتات ، عامل التركيز الحيوي ، مدينة كركوك.