

Original paper

Assessment of the Role of Gender in the Expression of UGT1A1 Gene in Hyperbilirubinemic Neonates

Hussein Hameed Hassan Mudhaji^{1*}, Dhaferah. J. A. AL-Fatlawi², Sabah N. AL-Fatlawi³

¹Faculty of physical planning, Univ. of kufa, AlNajaf, Iraq

²Faculty of education for girls, Univ. of kufa, AlNajaf, Iraq

³Najaf health directorate, Assadr hospital, unit of nephrology and kidney transplant, AlNajaf, Iraq

Abstract

Background: Neonatal jaundice (NJ) is a significant disease among neonates in Najaf province. It manifests 19% of the total deliveries and 50% of the in-patients, and affects about 60% of term and 80% of preterm neonates during the first week of life. UGT1A1 is a cause of concern in NJ, because it is the most important underlying cause of unconjugated hyperbilirubinemia.

Methods: A cohort of 85 neonate sorted into three groups according to the TSB level. Group 1 (TSB < 5 mg/dl), group 2 (TSB 5-15 mg/dl) and group 3 (TSB < 15 mg/l). UGT1A1 expression and TSB level was assessed for both males and females in each group.

Results & Discussion: Results showed high inverse correlation between UGT1A1 expression and the ratio of males ($r = -0.99$, $P < 0.01$), while females ratio expressed high direct correlation with UGT1A1 expression level with correlation coefficient of ($r = 0.99$, $P < 0.01$).

This might due to physiological maturity differences of liver between both sexes, and the plasma growth hormone (GH) pulses frequencies (released by the pituitary gland) which affects the expression of liver enzymes. It's found that GH pulses are higher in females than in males.

Conclusion: concluding that mean UGT1A1 expression level in females is more than that in the males during neonatal period therefore females are less vulnerable to NJ than males.

Aim: The aim of this study is to qualify the UGT1A1 gene expression in both males and females therefore determining health and health care requirements.

Keywords: Rule of gender in hyperbilirubinemia, Hyperbilirubinemia, Neonatal jaundice, UGT1A1 gene expression.

Introduction

Neonatal hyperbilirubinemia (NH) can be clinically defined as a yellowish discoloration of the patient's (Neonate's) sclera, skin and soft tissue as a result of bilirubin deposition ⁽¹⁾. It is first appears in the face and as the bilirubin level rises, it proceeds caudal to the trunk and then to the extremities which makes it appear on nail tips ⁽²⁻

⁴⁾. Neonatal jaundice (NJ) is a significant disease among neonates in Najaf province. According to Az-zahraa teaching hospital for pediatrics and obstetrics (ATHPO) 2012 records, Neonatal jaundice manifests 19% of the total deliveries and 50% of the in-patients. Internationally, NJ affect about 60% of term and 80% of preterm neonates during the first week of life ⁽⁵⁻⁷⁾. Neonatal jaundice is a cause of con-

*For Correspondence: E.mail hussein.abdulla@uokufa.edu.iq

cern for both parents and pediatricians⁽⁸⁾, because it is a threatening disease for the neonatal mental health and life, since its main complication is kernicterus (encephalopathy)⁽⁹⁻¹³⁾. Furthermore prevention of kernicterus is a time and money consuming process, It's found that prevention of one case of kernicterus in USA costing about (5.7) million dollars^(14,15), while in Ontario prevention one kernicterus case costing about (0.57) million dollars⁽¹⁶⁾. Unconjugated bilirubin is the prevalent in the neonatal hyperbilirubinemia⁽¹⁷⁾, this due to defect in either of the four mechanisms involved in bilirubin metabolism (bilirubin uptake, glucuronidation, excretion and entero-hepatic circulation), this defect either physiological which is mild and short term due to immaturity of neonate liver⁽¹⁸⁾ or pathological which is more sever, caused by either infection⁽¹⁹⁾ or genetic alterations in the UGT1A1 enzyme which represent the key element in the conjugation process^(20,21), or in the ligandins and membrane bound proteins of the hepatocytes membranes of the cell and organelle that responsible for bilirubin uptake⁽²²⁻²⁴⁾. Uridin diphosphate glucuronosyle transferase 1A1 (UGT1A1) is a cause

of concern in NJ, because it is the most important underlying cause of unconjugated hyperbilirubinemia⁽²⁵⁾ not only in neonatal period but along with the course of life. Impairment of UGT1A1 leads to impairment in detoxification of vast number of aromatic compounds including bilirubin, steroids (including steroid hormones) and some drugs such as irinotecan which is used as anti-cancer drug⁽²⁶⁻²⁸⁾. Data from Azzahraa Teaching Hospital for Obstetrics and Pediatrics revealed that NJ was more incident in males than females. UGT1A1 expression was examined to assess the the rule of gender in UGT1A1 expression, thereby aiding in individual and family disease management⁽²⁹⁻³¹⁾.

Methods

The present study was performed during the period from May to November 2013 in Az-zahraa Teaching Hospital for Pediatrics and Obstetrics and Annajaf Private Clinical Laboratory. The chemicals that used in this study listed in (tab. 1), whereas the apparatus and equipments used listed in (tab. 2).

Table 1.chemicals used in this study.

No.	Chemicals	Source
4	Grade water.	Parental- India
5	Human UGT1A1 ELISA kit	Cusabio- China

Table 2. Apparatus and Equipment used in this study.

No.	Equipment	Source
1	Centrifuge	Kokusan-19F Japan
2	ELISA reader	Bio-rad
3	Micropipette filtered tips.	Bioneer- Korea
4	Micropipette set (0.1-1000µl)	Biobasic- Canada
5	Multichannel micropipette	Ependrof
6	Timer alarm	China
7	U.V.-spectrophotometer	APEL- Japan
8	Incubator	Memmert- Germany
9	Refrigerator	Arjelic- Turkey

A cohort of 85 neonate subjects were used in this study, these cohorts sorted into three groups according to the TSB level. Group 1 represent the subjects whom TSB level was less than 5mg/dl, group 2 represent the subjects whom TSB level was from 5-15 mg/dl and group 3 represent the subjects whom TSB level was more than 15mg/l. UGT1A1 expression and TSB level was assessed for both males and females in each group.

A sample of 1ml venous blood was collected from these subjects, transferred to golden capped serum separating tubes (SST), left to clot for sufficient time, centrifuged at 3000 rpm for 15 min. Serum aliquot and stored in -24C° to the time the ELISA test where performed. TSB examined directly for the fresh sera. Reagents, standards and sera prepared for UGT1A1 assessment. UGT1A1 level assessed according to the manufacturer's protocol, optical density (OD) measured using ELISA reader and standard curve between OD and concentration; then drawn.

Curve expert software version (1.4) used to determine the standard curve fit and calculate the UGT1A1 concentration from analogous OD for each sample.

Results

Among groups statistics using student's t-test and correlation revealed that there is a high correlation between gender and UGT1A1 expression. The present study showed high inverse correlation between UGT1A1 expression and the ratio of males ($r = -0.99$, $P < 0.01$) (fig.1), while females ratio expressed high direct correlation with UGT1A1 expression level with correlation coefficient of ($r = 0.99$, $P < 0.01$) (fig. 2).

Discussion

Data analysis showed that mean UGT1A1 expression level in females is more than that in the males during neonatal period (tab. 3); so it is in accord with the results of ^{(32),(33)}. Therefore females are less vulnerable to severe NH than males. Liver-expressed genes are differing in males and females. Expression of these genes is affected by the profile of plasma growth hormone (GH) released by the pituitary gland in pulsatile form. The frequencies of GH pulses are higher in females than in males ⁽³⁴⁾.




Figure 1. UGT1A1 expression in male jaundiced neonates in the three groups.




Figure 2. UGT1A1 expression in female jaundiced neonates in the three groups.

Table 3. Relationship between gender and both UGT1A1 expression and TSB level, N=85.

Parameters	r ± SE	P-value
UGT1A1, M%	-0.99 ± 0.023	P<0.01
UGT1A1, F%	0.99 ± 0.023	P<0.01

Conclusion

Data analysis showed that mean UGT1A1 expression level in females is more than that in the males during neonatal period (tab. 1); therefore, females are less vulnerable to severe NH than males.

References

1. Madlon-Kay, D. J. (1997): Recognition of the Presence and Severity of Newborn Jaundice by Parents, Nurses, Physicians, and Ictermeter Pediatrics, Vol.100: e3.
2. Surhone, L. M.; Timpledon, M. T. and Marseken, S.F. (2010): Neonatal Jaundice. VDM Publishing, 6130514212, 9786130514211: 76.
3. Beeby P. and Evans N. (2011): Jaundice. RPA Newborn Care Guidelines Royal Prince Alfred Hospital, Vol.3.
4. Sampurna, K. and Shyamala, R. (2012): Hyperbilirubinemia in the term Newborn, J. ,
5. Piazza, A. J. and Stoll, B. J. (2008): Jaundice & hyperbilirubinaemia in the newborn, In Nelson Textbook of pediatrics. 18th ed. Philadelphia PA.. Saunders: 756-760.
6. NICE, clinical guideline (2010): National Collaborating Centre for Women's and Children's Health . clinical guideline, 2010: "Neonatal Jaundice". Royal college of Obstetricians and Gynaecologists: 33-52
7. Ferreira, D.; Oliveira, A. and Ferias, A. (2012): BMC Medical Informatics and Decision Making, Vol.12:143 (<http://www.biomedcentral.com/1472-6947/12/143>)..
8. Mishra, S.; Agrawal, R.; Deorari, A. K. and Paul, V. K. (2008): Jaundice in the newborns; Indian journal of pediatrics, 75: 157-163.
9. Petrova, A.; Mehta, R.; Birchwood, G.; Ostfeld, B. and Hegyi, T. (2006): Management of neonatal hyperbilirubinemia: Pediatricians' practices and educational needs; BMC Pediatrics, 6:6 doi:10.1186/1471-2431-6-6.
10. Burke, B. L.; Robbins, J. M.; Bird, T. M.; Hobbs, C. A. and Nesmith C. (2009): Trends in Hospitalizations for Neonatal Jaundice and Kernicterus in the United States, 1988 □2005; and John Mick Gilford; Pediatrics 2009:123-524.
11. Samar, N. El-Beshbishi; Karen, E. Shattuck; Amin, A. Mohammad and John, R. Petersen (2009): Hyperbilirubinemia and Transcutaneous Bilirubinometry Clinical Chemistry, 55:7.1280-1287 .
12. Gonçalves, A.; Costa, S.; Lopes, A.; Rocha, G.; Guedes, MB.; Centeno, MJ.; Silva, J.; Silva, MG.; Severo, M. and Guimarães, H., (2011):

- Prospective Validation of a Novel Strategy for Assessing Risk of Significant Hyperbilirubinemia. *Pediatrics*, Vol.127:e126. DOI: 10.1542/peds.2009-2771.
13. Burgos, A.E.; Flaherman V. J. and Newman T. B. (2012). Screening and Follow-Up for Neonatal Hyperbilirubinemia. A Review. *Clinical Pediatrics* 2012, Vol. 51: P 7–16.
 14. Gautham, K. Suresh and Robin, E. Clark, (2004): Cost-Effectiveness of Strategies That Are Intended to Prevent Kernicterus in Newborn Infants. *Pediatrics*, 114: 917-924.
 15. Maisels, M. J.; Vinod, K. B.; Bogen D.; Thomas, B. N.; Ann, R. S. and Jon F. W. (2009) : Hyperbilirubinemia in the Newborn Infant 35 Weeks' Gestation. An Update With Clarifications. *Pediatrics*, Vol.124(1193). DOI: 10.1542/peds.2009-0329
 16. Xie, B.; et al, (2012): Cost-effectiveness analysis of a system-based approach for managing neonatal jaundice and preventing kernicterus in Ontario. *Paediatr Child Health*, 17:11-16.
 17. Gilmour, S.M.(2004): Prolonged neonatal jaundice. When to worry and what to do. *Paediatr Child Health*, Vol.9: P700-704.
 18. Mladenović, M.; Radlović, N.; Ristić, D.; Leković, Z.; Radlović, P.; Pavlović, M.; Gajić, M.; Puskarević, M.; Davidović, I. and Djurdjević, J.(2007): Arias icterus--prolonged unconjugated hyperbilirubinemia caused by breast milk. *SrpArhCelok Lek.*, 135. 11-12:655-8.
 19. Smith, P. A.; Larenas, J.; Contreras, J.; Cassigoli, J.; Venegas, C.; Rojas, M. E.; Guajardo, A.; Pérez, S. and Díaz S.(2006): Infectious haemolytic anaemia causes jaundice outbreaks in seawater-cultured coho salmon, *Oncorhynchus kisutch* (Walbaum), in Chile.; *J Fish Dis.*, 29:709-15.
 20. Tiribelli, C. and Donald, O. J.(1996): New Concepts in Bilirubin and Jaundice: Report of the Third International Bilirubin Workshop, April 6-8, 1995, Trieste, Italy. *hepatology* Vol. 24: 1297.
 21. Buonocore, G., Bracci, R. and Weindling, M.(2012): Bilirubin metabolism; Neonatology, A practical approach to neonatal management, Ch. 81:608.
 22. Stephen, D. Z. and Goessling, W.(2000): Mechanism of hepatocellular uptake of albumin-bound bilirubin. *Biochimica et Biophysica Acta*, 1463: 197-208.
 23. Wang, X.; Chowdhury, J. R. and Chowdhury N. R. (2006): Bilirubin metabolism: Applied physiology; Elsevier Current Paediatrics, 16: 70–74.
 24. Michael D. C.(2010): Human drug metabolism an introduction; Aston University. Birmingham. UK. A John Wiley & Sons, Ltd., Publication, 2ed: 125-129.
 25. Servedio, V.; Apolito, M.; Maiorano, N.; Minuti, B.; Torricelli, F.; Ronchi, F.; Zancan, L.; Perrotta, S.; Vajro, P.; Boschetto, L. and Iolascon, A.(2005): Spectrum of UGT1A1 Mutations in Crigler-Najjar (CN) Syndrome Patients: Identification of Twelve Novel Alleles and Genotype-Phenotype Correlation; MUTATION IN BRIEF; WILEY-LISS, INC.; #793: DOI: 10.1002/humu.9322.
 26. Iyer, L.; et al(1998): Genetic predisposition to the metabolism of irinotecan (CPT-11). Role of uridine diphosphate glucuronosyl transferase isoform 1A1 in the glucuronidation of its active metabolite (SN-38) in human liver microsomes. *J Clin Invest*, 101:847-54.
 27. Beutler, E.; Gelbart, T., Demina, A. (1998): Racial variability in the UDP-glucuronosyltransferase 1 (UGT1A1) promoter. A balanced polymorphism for regulation of bilirubin metabolism. *Proc Natl Acad Sci U S A.*, Vol.95 :8170-4.
 28. Ramírez, J.; Ratain, M. J. and Innocenti, F.(2010): Uridine 5'-diphosphoglucuronosyltransferase genetic polymorphisms and response to cancer chemotherapy. *Future Oncol.* , 6: 563–585., doi:10.2217/fon.10.17.
 29. Chen, M. H.; Su, Y. N.; Hsieh, W. S.; Chou, H. C.; Chen, C. Y. and Tao, P. N.(2008): UDP-Glucuronosyl Transferase 1A1 (UGT1A1) Gene Polymorphism in Neonatal Hyperbilirubinemia –Preliminary Report. *Clinical Neonatology*, Vol. 15: P 20.
 30. Skierka, J. M.; Kotzer, K. E; Lagerstedt, S. A.; O'Kane, D. J. and Baudhuin, L. M.(2013): UGT1A1 genetic analysis as a diagnostic aid for individuals with unconjugated hyperbilirubinemia. *J Pediatr.* , 162: 1146-52, 1152.e1-2. doi: 10.1016/j.jpeds. 2012.11.042. Epub 2013 Jan 4.
 31. Payton, J. and Lockwood, C.(2013): UGT1A1-28 Genotyping for Irinotecan Toxicity; Molecular Diagnostic Laboratory; Washington university in st. Louis school of medicin, <http://pathology.wustl.edu/patientcare/molldiagnostic.ph>.

32. Chang, J.L.; et al, (2007): *UGT1A1* Polymorphism Is Associated with Serum Bilirubin Concentrations in a Randomized, Controlled, Fruit and Vegetable Feeding Trial. American Society for Nutrition, 137 : 890-897.
33. Zisk, J.L., et al(2011): Do premature female infants really do better than their male counterparts?.Am J Perinatol., 28:241-6.
34. David, J. and Minita, G.(2009): Sex Differences in the Expression of Hepatic Drug Metabolizing Enzymes. MolPharmacol 76:215–228.